

On Ruelle’s Lemma and Ruelle Zeta Functions

Paul Wright

January 19, 2014

Abstract

In this article we prove an important inequality regarding the Ruelle operator in hyperbolic flows. This was already proven briefly by Mark Pollicott and Richard Sharp in a low dimensional case [PS1], but we present a clearer proof of the inequality, filling in gaps and explaining the ideas in more detail, and extend the inequality to higher dimensional flows. This inequality is necessary to prove a proposition about the analyticity of Ruelle zeta functions.¹

1 Introduction

This article gives a comprehensive and rigorous proof of a lemma by Mark Pollicott and Richard Sharp (called “Lemma 2” in [PS1]). They claim that the result was essentially proved in a paper by David Ruelle [Ru], but Ruelle’s paper does not state or prove the lemma explicitly. Another proof of the lemma can be found in a paper by Frédéric Naud [Na], which clarifies some issues with the proof in [PS1] but contains a new error. A similar but weaker result was also proven in [PS2]. The three papers [Na, PS1, PS2] each prove the lemma briefly for low dimensional flows but each proof has significant gaps and even errors. In this article the proof follows Naud’s approach more closely than the original Pollicott-Sharp proof. We recover the original result from [PS1], but fill in gaps in the proof and extend the result to higher dimensional flows.

Let M be a compact smooth manifold and let $\phi_t : M \rightarrow M$ be an Axiom A flow [PP1]. Then M contains a non-wandering set that can be decomposed into basic sets $\Lambda_i \subset M$ on which ϕ_t is hyperbolic and transitive, its periodic points are dense and there exists a neighbourhood $U \supset \Lambda_i$ with $\Lambda_i = \bigcap_{n=-\infty}^{\infty} f^n(U)$. See [PP1] for full definitions. We use the fact that the periodic points are dense in Λ near the end of the paper. We focus on a single basic set Λ .

For any $x \in M$ let $W_\varepsilon^s(x), W_\varepsilon^u(x)$ be the local stable and unstable manifolds through x respectively, as defined in [PP1]. We call a subset A of a stable or unstable manifold *admissible* if $A = \overline{\text{Int } A}$, where the closure and interior are in the induced topology of the local stable or unstable manifold intersected with Λ .

¹This article represents the main result of the author’s honours dissertation at the University of Western Australia, supervised by Luchezar Stoyanov.

A *Markov rectangle* R_i can be constructed from a point $z_i \in \Lambda$ and admissible subsets $U_i \subset W_\varepsilon^u(z_i) \cap \Lambda$ and $S_i \subset W_\varepsilon^s(z_i) \cap \Lambda$ using a local product structure on Λ [Do1, PP1]. The interior of a rectangle $\text{Int } R_i$ can be similarly constructed from $\text{Int } U_i$ and $\text{Int } S_i$. A Markov family of rectangles $R_1, \dots, R_k \subset \Lambda$ can be constructed as described in [Do1, PP1], and we let $R = \bigcup_{i=1}^k R_i$.

In [PS1], the stable and unstable leaves are always one-dimensional and each U_i is called an interval. We are working with higher dimensional flows, so the unstable leaves could in general have a higher dimension. We call each U_i a *Markov leaf*. The set of Markov leaves is disjoint. We denote $U = \bigcup_{i=1}^k U_i$ and $\text{Int } U = \bigcup_{i=1}^k \text{Int } U_i$.

Define the Poincaré map $H : R \rightarrow R$ by $H(x) = \phi_{r(x)}(x) \in R$, where $r(x) > 0$ is the *first return time*, the smallest positive time such that $\phi_{r(x)}(x) \in R$.

Define a $k \times k$ matrix A by

$$A(i, j) = \begin{cases} 1 & : H(\text{Int } R_i) \cap \text{Int } R_j \neq \emptyset \\ 0 & : \text{otherwise.} \end{cases}$$

We use this matrix to define the symbol spaces X_A and X_A^+

$$\begin{aligned} X_A &= \{\xi = \{\xi_n\}_{n=-\infty}^{\infty} : \xi_n \in \{1, \dots, k\}, A(\xi_n, \xi_{n+1}) = 1, \forall n \in \mathbb{Z}\} \\ X_A^+ &= \{x = \{\xi_n\}_{n=0}^{\infty} : \xi_n \in \{1, \dots, k\}, A(\xi_n, \xi_{n+1}) = 1, \forall n \geq 0\}. \end{aligned}$$

It is always possible to construct the Markov family such that the matrix A is *irreducible* and *aperiodic* [PP1].

Define the *stable holonomy map* $\pi^u : R \rightarrow U$ by $\pi^u(x) = W_\varepsilon^s(x) \cap U_i$. Define the expanding map $f : U \rightarrow U$ by $f = \pi^u \circ H|_U$. This map is expanding in the sense that there exist constants $0 < \gamma < 1$ and $C_1 > 0$ such that, if $f^j(x)$ and $f^j(y)$ are on the same Markov leaf U_{i_j} for every $0 \leq j \leq m$, we have

$$d(x, y) \leq C_1 \gamma^m d(f^m x, f^m y). \quad (1)$$

2 Preliminaries

2.1 Cylinders

Definition 2.1. For each $n \geq 0$, consider all strings of length n of symbols $1, 2, \dots, k$. We call a string $\alpha = (\alpha_0, \dots, \alpha_{n-1})$ a *word* if it is admissible under A , that is, if $A(\alpha_j, \alpha_{j+1}) = 1$ for all $0 \leq j \leq n-2$. For any word α we write $|\alpha| = n$ and define a subset of one of the Markov leaves

$$U_\alpha = U_{\alpha_0} \cap f^{-1}U_{\alpha_1} \cap \dots \cap f^{-(n-1)}U_{\alpha_{n-1}}.$$

Here f^{-1} denotes the preimage. We call this subset a *cylinder* of length n in the leaf U_{α_0} . Each U_i is a cylinder of length 1 and is always a compact set, but other cylinders can be open, closed, or neither. If $|\alpha| = |\beta| > 1$ then either $\alpha = \beta$ or $U_\alpha \cap U_\beta = \emptyset$. This is not true in general for the closures $\overline{U_\alpha}$ and $\overline{U_\beta}$. A cylinder U_α is always nonempty, and as the length of a string approaches infinity, the corresponding cylinder approaches a single point. Thus a single point in U can be represented by an infinite string of symbols, i.e. an element of X_A^+ .

Definitions 2.2. Let $\alpha = (\alpha_0, \dots, \alpha_{n-1})$ be a word of length n and let $i = 1, \dots, k$. Then if $A(\alpha_{n-1}, i) = 1$ and $A(i, \alpha_0) = 1$, we define

$$\alpha i = (\alpha_0, \dots, \alpha_{n-1}, i), \quad i\alpha = (i, \alpha_0, \dots, \alpha_{n-1}), \quad \text{and } \bar{\alpha} = (\alpha_0, \dots, \alpha_{n-2}).$$

In general the expanding map f^n is not always injective for $n \geq 1$. For each permissible word α we define an inverse that follows the word backwards. This is always possible for points on the interior of a leaf. Let α be a word of length n and let $x \in f(U_{\alpha_{n-1}}) \cap \text{Int } U_i$ for any i with $A(\alpha_{n-1}, i) = 1$. Define $f_\alpha^{-1}(x)$ to be the unique point y such that $f^n(y) = x$ and $y \in U_\alpha$. For a single symbol i we can write $ix = f_i^{-1}(x)$.

2.2 Hölder functions

Definition 2.3. For any $A \subseteq U$, denote by $C(A)$ the set of all continuous functions $w : A \rightarrow \mathbb{C}$, and let $w \in C(A)$. For any $0 < \mu \leq 1$, we say that w is μ -Hölder on A , or Hölder continuous on A with exponent μ , if there exists $C_2 > 0$ such that $|w(x) - w(y)| \leq C_2 d(x, y)^\mu$ for all $x, y \in A \cap U_i$, $i = 1, \dots, k$. We define the μ norm of w on A by

$$|w|_\mu = \sup \left\{ \frac{|w(x) - w(y)|}{d(x, y)^\mu} : x, y \in A \cap U_i, i \in \{1, \dots, k\}, x \neq y \right\}.$$

When $\mu = 1$ we say w is *Lipschitz* or *Lipschitz continuous*. For any admissible set A , we denote by $H^\mu(A)$ the set of all functions that are μ -Hölder on A . That is,

$$H^\mu(A) = \{w \in C(U) : |w|_\mu < \infty \text{ on } A\}.$$

We also define the infinity norm $|w|_\infty = \sup_{x \in U} |w(x)|$ and a total norm

$$\|w\|_\mu = |w|_\mu + |w|_\infty.$$

For an operator $L : C(U) \rightarrow C(U)$ we use the operator norm

$$\|L\|_\mu = \sup \left\{ \frac{\|Lw\|_\mu}{\|w\|_\mu} : w \in C(U), w \neq 0 \right\}.$$

Pollicott and Sharp use a different (C^1) norm in their paper [PS1].

If $f : A \rightarrow \mathbb{C}$ is μ -Hölder on some set A in any metric space, then there exists a unique extension $\text{ext}(f) : \bar{A} \rightarrow \mathbb{C}$ of f to the closure of A , such that $\text{ext}(f)$ is μ -Hölder on \bar{A} [WW]. We call $\text{ext}(f)$ the *Hölder extension* of f .

Lemma 2.4. *If w is μ -Hölder on a set containing x and y , then*

$$\begin{aligned} |e^{w(x)-w(y)} - 1| &\leq |w(x) - w(y)| e^{|w(x)-w(y)|} \\ &\leq |w|_\mu e^{|w|_\mu |U|^\mu} d(x, y)^\mu. \end{aligned}$$

2.3 Characteristic Function

For a word α of length n we let χ_α denote the characteristic function

$$\chi_\alpha(x) = \chi_{U_\alpha}(x) = \begin{cases} 1 & \text{if } x \in U_\alpha, \\ 0 & \text{otherwise.} \end{cases}$$

Observe that for any word α of length n , $\chi_\alpha(x)$ is not Lipschitz on all U_i , but it is piecewise Lipschitz. In particular, it is Lipschitz (and hence μ -Hölder) on U_β for any word β with $|\beta| \geq |\alpha|$, with Lipschitz and Hölder constants $|\chi_\alpha(x)|_{\text{Lip}} = |\chi_\alpha(x)|_\mu = 0$. We write

$$\chi_\alpha \in \bigoplus_{|\beta|=n} H^\mu(U_\beta).$$

2.4 First Return Map

For $x \in U$, we write $r^n(x) = r(x) + r(fx) + \dots + r(f^{n-1}x)$. In a three dimensional flow, the first return map r is always Lipschitz continuous on 2-cylinders $U_{i,j}$ for any $i, j \in 1, \dots, k$ with $A(i, j) = 1$. However in higher dimensions r is not Lipschitz continuous in general, but it is always Hölder continuous on each $U_{i,j}$ [Ha]. Henceforth μ refers to the largest exponent such that r is μ -Hölder on every $U_{i,j}$. Then $|r(x)| \leq |r|_\infty$ and $|r(x) - r(y)| \leq |r|_\mu d(x, y)^\mu$ for any x, y in the same 2-cylinder.

Lemma 2.5. *Whenever x and y are on the same cylinder U_α with $|\alpha| = m$, there exists a constant B_1 depending only on r, C_1, γ and μ such that*

$$|r^m(x) - r^m(y)| \leq B_1 d(f^{m-1}x, f^{m-1}y)^\mu.$$

Proof.

$$\begin{aligned} |r^m(x) - r^m(y)| &\leq \sum_{j=0}^{m-1} |r(f^j x) - r(f^j y)| \leq \sum_{j=0}^{m-1} |r|_\mu d(f^j x, f^j y)^\mu \\ &\leq \sum_{j=0}^{m-1} |r|_\mu (\gamma^{m-j-1} C_1 d(f^{m-1}x, f^{m-1}y))^\mu \\ &\leq |r|_\mu C_1^\mu d(f^{m-1}x, f^{m-1}y)^\mu \sum_{i=0}^{m-1} (\gamma^\mu)^i \\ &\leq |r|_\mu C_1^\mu \frac{1}{1 - \gamma^\mu} d(f^{m-1}x, f^{m-1}y)^\mu \\ &= B_1 d(f^{m-1}x, f^{m-1}y)^\mu \end{aligned}$$

□

2.5 The transfer operator and zeta function

Let $s = a + ib \in \mathbb{C}$, with $|a| \leq a_0$ and $|b| \geq b_0$ for some constants a_0, b_0 . Let h denote the *entropy* of the flow ϕ_t , and for any continuous $w : U \rightarrow \mathbb{R}$ let $P(w)$ denote the *pressure* of w . Definitions of these can be found in [W]. We have $P(0) = h$.

Let $\hat{U} \subset U$ be the set of points $x \in \text{Int } U$ such that $f^j x \in \text{Int } U$ for any integer $j \geq 0$. We define the *Ruelle transfer operator* $L_{-sr} : C(\hat{U}) \rightarrow C(\hat{U})$ by

$$L_{-sr}w(x) = \sum_{fy=x} e^{-sr(y)}w(y).$$

When $x \notin \hat{U}$, r may not be continuous at y for $fy = x$, and hence $L_{-sr}w$ may not be continuous at x . When $L_{-sr}w$ is Hölder continuous on \hat{U} , we extend this function to points in U using the Hölder extension, by defining $L_{-sr}w(x) = \text{ext}(L_{-sr}w)(x)$.

Define the zeta function $\zeta(s)$ for the flow ϕ_t by

$$\zeta(s) = \prod_{\tau} (1 - e^{-sl(\tau)})^{-1},$$

where the product is over prime periodic orbits τ , i.e. $f^n(x) = x, \forall x \in \tau$ for some n , and $l(\tau) = r^n(x)$ for some $x \in \tau$ is the length of the orbit. This zeta function converges to a nonzero analytic function for $\Re(s) > h$, where h is the entropy of the flow ϕ_t [PS1]. It is important to choose a_0 large enough that the compact interval $I = [-a_0, a_0]$ contains h .

For all $s \in \mathbb{C}$, we define a weighted sum on the periodic points

$$Z_n(-sr) = \sum_{f^n(x)=x} e^{-sr^n(x)}.$$

We can use the fact that

$$\zeta(s) = \exp \sum_{n=1}^{\infty} \frac{1}{n} Z_n(-sr)$$

(see [PS1]) to relate the zeta function to the Ruelle transfer operator L_{-sr} , using Ruelle's Lemma, which relates $Z_n(-sr)$ to the Ruelle operator L_{-sr} .

3 Ruelle's Lemma

Theorem 3.1. (*Ruelle's Lemma*) *For each Markov leaf U_i , fix an arbitrary point $x_i \in U_i$. For all $\varepsilon > 0, a_0 > 0$ and $b_0 > 0$, there exists $C_\varepsilon > 0$ such that*

$$\left| Z_n(-sr) - \sum_{i=1}^k L_{-sr}^n \chi_i(x_i) \right| \leq C_\varepsilon |\Im(s)| \sum_{m=2}^n \left(\|L_{-sr}^{n-m}\|_{\mu} (\gamma^\mu)^m e^{m(\varepsilon + P(-ar))} \right),$$

for all $s = a + ib \in \mathbb{C}$ with $|a| \leq a_0$ and $|b| \geq b_0$. Here μ denotes the Hölder exponent of r and $P(-ar)$ denotes the pressure of $-ar$.

Our aim in this article is to give a rigorous proof of this lemma.

3.1 Proof

Fix an arbitrary point $x_i \in U_i$ for each Markov leaf U_i . Let $\varepsilon > 0$, $a_0 > 0$ and $b_0 > 0$.

Proposition 3.2. *Let $m \geq 1$. Given a function w that is μ -Hölder on all cylinders of length $m + 1$, the transfer operator of w is always μ -Hölder on all m -cylinders. That is,*

$$L_{-sr} : \bigoplus_{|\alpha|=m+1} H^\mu(U_\alpha) \rightarrow \bigoplus_{|\alpha|=m} H^\mu(U_\alpha).$$

Proof. It suffices to show that for any particular α , if w is μ -Hölder on $U_{i\alpha}$ for all i with $A(i, \alpha_0) = 1$, then $L_{-sr}w$ must be μ -Hölder on $\text{Int } U_\alpha$. Let $w \in H^\mu(U_{i\alpha})$ and let $x, y \in \text{Int } U_\alpha$. Then using Lemma 2.4, we have

$$\begin{aligned} |L_{-sr}w(x) - L_{-sr}w(y)| &= \left| \sum_{A(i, \alpha_0)=1} e^{-sr(ix)} w(ix) - \sum_{A(i, \alpha_0)=1} e^{-sr(iy)} w(iy) \right| \\ &\leq \sum_{A(i, \alpha_0)=1} |e^{-sr(iy)}| \left(|e^{sr(iy)-sr(ix)} - 1| |w(ix)| + |w(iy) - w(ix)| \right) \\ &\leq \sum_{A(i, \alpha_0)=1} |e^{-sr(iy)}| (|s||r|_\mu e^{a_0|r|_\mu |U|^\mu} d(ix, iy)^\mu |w(ix)| + |w|_\mu d(ix, iy)^\mu) \\ &\leq ke^{a_0|r|_\infty} (|s||r|_\mu e^{a_0|r|_\mu |U|^\mu} |w|_\infty + |w|_\mu) C_1^\mu \gamma^\mu d(x, y)^\mu \\ &\leq C_4 \|w\|_\mu d(x, y)^\mu, \end{aligned}$$

for some constant C_4 . So $L_{-sr}w$ is μ -Hölder on $\text{Int } U_\alpha$. Using the Hölder extension, $L_{-sr}w$ is μ -Hölder on every U_α . □

Corollary 3.3. *In particular, if $|\alpha| = m$ and $n \geq m$ then $L_{-sr}^n \chi_\alpha \in H^\mu(U_i)$ for all $i = 1, \dots, k$.*

Since the map f is expanding, any cylinder U_α can only contain zero or one n -periodic points, that is, a point $x \in U_\alpha$ such that $f^n x = x$ where $n = |\alpha|$. For each string α , we have to choose a point $x_\alpha \in U_\alpha$ in a particular way. We choose a point from the cylinder as follows:

- (1) if U_α has an n -periodic point, then let $x_\alpha \in U_\alpha$ be such that $f^n x_\alpha = x_\alpha$;
- (2) if U_α has no n -periodic point and $n > 1$, then choose $x_\alpha \in U_\alpha$ arbitrarily such that $x_\alpha \notin f(U_{\alpha_{n-1}})$;
- (3) if $|\alpha| = n = 1$, then U_α has no n -periodic point. Choose $x_\alpha = x_i$ (where $i = \alpha_0$ and $x_i \in U_i$ is one of the points we fixed from the start).

Part of the second condition was introduced by Naud [Na]. Both [Na] and [PS1] leave out the last condition for $|\alpha| = 1$, but [PS2] includes it. The need for this condition becomes clear in (2). Note that this is consistent with the other two, that is if $n = 1$ and we choose $x_\alpha = x_i$, we still have that $x_\alpha \notin f(U_{\alpha_{n-1}})$ and x_α is not a 1-periodic point (i.e. a fixed point). This is because $A(i, i) = 0$ for all i , so there are no fixed points.

Lemma 3.4. *By choosing x_α in this way, we have*

$$(L_{-sr}^n \chi_\alpha)(x_\alpha) = \begin{cases} e^{-sr^n(x_\alpha)} & \text{if } x_\alpha \text{ is a periodic point,} \\ 0 & \text{otherwise.} \end{cases}$$

Proof. We have

$$(L_{-sr}^n \chi_\alpha)(x_\alpha) = \sum_{f^n y = x_\alpha} e^{-sr^n(y)} \chi_\alpha(y),$$

so we are looking for points y such that $f^n y = x_\alpha$ and $y \in U_\alpha$.

Suppose U_α has an n -periodic point. Then there is exactly one point y satisfying both $f^n y = x_\alpha$ and $\chi_\alpha(y) = 1$. Only x_α satisfies both of these, so we have $y = x_\alpha$. Thus $(L_{-sr}^n \chi_\alpha)(x_\alpha) = e^{-sr^n(x_\alpha)}$ when U_α has an n -periodic point.

Suppose U_α has no n -periodic point. Then $x_\alpha \in U_\alpha$ but $x_\alpha \notin f(U_{\alpha_{n-1}})$. Suppose for a contradiction that there is some $y \in U_\alpha$ s.t. $f^n y = x_\alpha$. Then

$$y \in U_\alpha \Rightarrow y \in f^{-(n-1)} U_{\alpha_{n-1}} \Rightarrow f^n y \in f(U_{\alpha_{n-1}}) \Rightarrow x_\alpha \in f(U_{\alpha_{n-1}}).$$

This is a contradiction, so the sum is empty. Thus $(L_{-sr}^n \chi_\alpha)(x_\alpha) = 0$ when U_α has no periodic point. □

Lemma 3.5.

$$Z_n(-sr) = \sum_{|\alpha|=n} (L_{-sr}^n \chi_\alpha)(x_\alpha)$$

Proof. Recall that $Z_n(-sr)$ is a sum over all period n points of U . Since $U = \bigcup_{|\alpha|=n} U_\alpha$, we can break up this sum as follows

$$Z_n(-sr) = \sum_{f^n x = x} e^{-sr^n(x)} = \sum_{|\alpha|=n} \sum_{f^n x = x} e^{-sr^n(x)} \chi_\alpha(x).$$

Suppose U_α has an n -periodic point. Then it is equal to x_α , so we have

$$\sum_{f^n x = x} e^{-sr^n(x)} \chi_\alpha(x) = e^{-sr^n(x_\alpha)} = (L_{-sr}^n \chi_\alpha)(x_\alpha).$$

Suppose U_α has no n -periodic points. Then we have

$$\sum_{f^n x = x} e^{-sr^n(x)} \chi_\alpha(x) = 0 = (L_{-sr}^n \chi_\alpha)(x_\alpha).$$

Then,

$$Z_n(-sr) = \sum_{|\alpha|=n} (L_{-sr}^n \chi_\alpha)(x_\alpha).$$

□

Note that, with the third condition we placed on x_α , it is trivial to show that

$$\sum_{i=1}^k (L_{-sr}^n \chi_{U_i})(x_i) = \sum_{|\alpha|=1} (L_{-sr}^n \chi_\alpha)(x_\alpha). \quad (2)$$

Proposition 3.6.

$$Z_n(-sr) - \sum_{i=1}^k L_{-sr}^n \chi_i(x_i) = \sum_{m=2}^n \left(\sum_{|\alpha|=m} L_{-sr}^n \chi_\alpha(x_\alpha) - \sum_{|\beta|=m-1} L_{-sr}^n \chi_\beta(x_\beta) \right)$$

Proof. Expand the sum on the RHS and notice that almost all the terms cancel out, leaving only

$$\sum_{|\alpha|=n} (L_{-sr}^n \chi_\alpha)(x_\alpha) - \sum_{|\alpha|=1} (L_{-sr}^n \chi_{U_\alpha})(x_\alpha).$$

Using Lemma 3.5 and (2), this is equal to

$$Z_n(-sr) - \sum_{i=1}^k L_{-sr}^n \chi_i(x_i).$$

□

Note that for any word β of length $m-1$, we have $U_\beta = \bigcup_{A(\beta_{m-2}, i)=1} U_{\beta i}$, so for any $x \in M$ we can write

$$\chi_\beta(x) = \sum_{A(\beta_{m-2}, i)=1} \chi_{\beta i}(x).$$

So we have, for $m \leq n$,

$$\sum_{|\beta|=m-1} L_{-sr}^n \chi_\beta(x_\beta) = \sum_{|\beta|=m-1} \sum_{A(\beta_{m-2}, i)=1} L_{-sr}^n \chi_{\beta i}(x_\beta) = \sum_{|\alpha|=m} L_{-sr}^n \chi_\alpha(x_{\bar{\alpha}}),$$

and therefore

$$Z_n(-sr) - \sum_{i=1}^k L_{-sr}^n \chi_i(x_i) = \sum_{m=2}^n \sum_{|\alpha|=m} [L_{-sr}^n \chi_\alpha(x_\alpha) - L_{-sr}^n \chi_\alpha(x_{\bar{\alpha}})].$$

By Corollary 3.3, $L_{-sr}^n \chi_\alpha$ is μ -Hölder everywhere on U . So we have, for all $n \geq 2$,

$$\begin{aligned} |L_{-sr}^n \chi_\alpha(x_\alpha) - L_{-sr}^n \chi_\alpha(x_{\bar{\alpha}})| &\leq \|L_{-sr}^n \chi_\alpha\|_\mu d(x_\alpha, x_{\bar{\alpha}})^\mu \\ &\leq \|L_{-sr}^{n-m}\|_\mu \cdot \|L_{-sr}^m \chi_\alpha\|_\mu d(x_\alpha, x_{\bar{\alpha}})^\mu, \end{aligned}$$

where $\|\cdot\|_\mu$ is the operator norm derived from the μ -Hölder norm.

So we have the following estimate for all $n \geq 2$:

$$\left| Z_n(-sr) - \sum_{i=1}^k L_{-sr}^n \chi_i(x_i) \right| \leq \sum_{m=2}^n \sum_{|\alpha|=m} \|L_{-sr}^{n-m}\|_\mu \cdot \|L_{-sr}^m \chi_\alpha\|_\mu d(x_\alpha, x_{\bar{\alpha}}). \quad (3)$$

3.2 Particular estimates

The last part of the proof is to find particular estimates for the three parts in the RHS of equation 3.

1. In the low dimensional case, $\mu = 1$, so $\|L_{-sr}^{n-m}\|_1$ can sometimes be estimated using an important theorem of Dolgopyat [PS1]. In this proof we leave it as $\|L_{-sr}^{n-m}\|_\mu$. Later we will show an example of how Dolgopyat's theorem can be used in some cases along with the lemma, to get an extension of the zeta function.
2. $\|L_{-sr}^m \chi_\alpha\|_\mu$ is simply estimated by $\|L_{-sr}^m \chi_\alpha\|_\mu \leq C|\Im(s)|$ in the Pollicott-Sharp paper [PS1]. Naud produces a more detailed estimate that we cover in detail in the next section.
3. Since $x_\alpha, x_{\bar{\alpha}}$ are in the same cylinder $U_{\bar{\alpha}}$, (1) gives us

$$d(x_\alpha, x_{\bar{\alpha}})^\mu \leq C_1^\mu \gamma^{\mu(m-2)} d(f^{m-2} x_\alpha, f^{m-2} x_{\bar{\alpha}})^\mu \leq C_5 (\gamma^\mu)^m, \quad (4)$$

for some constant $C_5 > 0$.

We now fix $m \geq 1$ and attempt to estimate $\|L_{-sr}^m(\chi_\beta)\|_\mu$.

Lemma 3.7.

$$(L_{-sr}^m \chi_\beta)(x) = \begin{cases} e^{-r^m(f_\beta^{-1}x)} & \text{if } x \in f(U_{\beta_{m-1}}), \\ 0 & \text{otherwise.} \end{cases}$$

Proof. We have $(L_{-sr}^m \chi_\beta)(x) = \sum_{f^m y = x} e^{-sr^m(y)} \chi_\beta(y)$.

Suppose $x \in f(U_{\beta_{m-1}})$. Then $f^m y = x$ and $\chi_\beta(y) = 1$ implies $y = f_\beta^{-1}(x)$.

So there is only one non-zero term in this sum, which is $e^{-r^m(f_\beta^{-1}x)}$.

Now suppose $x \notin f(U_{\beta_{m-1}})$. Then if $f^m y = x$ and $\chi_\beta(y) = 1$, then $f^{m-1}y \in U_{\beta_{m-1}}$, so $x = f^m(y) \in f(U_{\beta_{m-1}})$, which is a contradiction. So the sum is zero. \square

For each admissible word β with $|\beta| = m$, we fix a point $y_\beta \in f(U_{\beta_{m-1}})$. We will see later how to choose y_β . Set $z_\beta = f_\beta^{-1}(y_\beta)$.

Lemma 3.8.

$$\|L_{-sr}^m(\chi_\beta)\|_\mu \leq (e^{a_0|U|^\mu B_1} + B_1|s|e^{a_0|U|^\mu(1+\gamma^\mu)B_1})e^{-ar^m(z_\beta)}.$$

Proof. Let $|U| = \max_i \text{diam}(U_i)$ be the largest diameter possible for a Markov leaf. Let x, y be in the same Markov leaf. If $x, y \notin f(U_{\beta_{m-1}})$, then clearly $|L_{-sr}^m(\chi_\beta)(x)| = |L_{-sr}^m(\chi_\beta)(x) - L_{-sr}^m(\chi_\beta)(y)| = 0$. Otherwise, both x and y are in $f(U_{\beta_{m-1}})$.

x and y_β may not be on the same leaf, but $f_\beta^{-1}x$ and $f_\beta^{-1}y_\beta$ are on the same cylinder U_β . So using Lemmas 3.7 and 2.5, we have

$$\begin{aligned} |L_{-sr}^m\chi_\beta(x)| &= |e^{-(a+ib)r^m(f_\beta^{-1}x)}| = e^{-ar^m(f_\beta^{-1}x)} \\ &\leq e^{|a||r^m(f_\beta^{-1}x) - r^m(f_\beta^{-1}y_\beta)|} e^{-ar^m(f_\beta^{-1}y_\beta)} \\ &\leq e^{a_0B_1d(f_\beta^{-1}x, f_\beta^{-1}y_\beta)^\mu} e^{-ar^m(z_\beta)} \leq e^{a_0|U|^\mu B_1} e^{-ar^m(z_\beta)}. \end{aligned}$$

Since $f_\beta^{-1}x$ and $f_\beta^{-1}y$ are on the same cylinder U_β , and x, y are on the same leaf, we have $|r^m(f_\beta^{-1}x) - r^m(f_\beta^{-1}y)| \leq \gamma^\mu B_1 d(x, y)$. Using Lemma 2.4, we get

$$\begin{aligned} |L_{-sr}^m(\chi_\beta)(x) - L_{-sr}^m(\chi_\beta)(y)| &\leq |e^{-sr^m(f_\beta^{-1}x) + sr^m(f_\beta^{-1}y)} - 1| |e^{-sr^m(f_\beta^{-1}y)}| \\ &\leq \left(|s| B_1 \gamma^\mu d(x, y)^\mu e^{|a|B_1\gamma^\mu|U|^\mu} \right) \left(e^{a_0B_1|U|^\mu} e^{-ar^m(z_\beta)} \right) \\ &\leq B_1 \gamma^\mu |s| e^{a_0B_1(\gamma^\mu+1)|U|^\mu} e^{-ar^m(z_\beta)} d(x, y)^\mu. \end{aligned}$$

Combining these results we get

$$\begin{aligned} \|L_{-sr}^m(\chi_\beta)\|_\mu &= |L_{-sr}^m(\chi_\beta)|_\infty + |L_{-sr}^m(\chi_\beta)|_\mu \\ &\leq e^{a_0|U|^\mu B_1} e^{-ar^m(z_\beta)} + B_1 \gamma^\mu |s| e^{a_0|U|^\mu(1+\gamma^\mu)B_1} e^{-ar^m(z_\beta)} \\ &\leq (e^{a_0|U|^\mu B_1} + B_1 \gamma^\mu |s| e^{a_0|U|^\mu(1+\gamma^\mu)B_1}) e^{-ar^m(z_\beta)}. \end{aligned}$$

□

Now $|s| = |a + bi| \leq |a| + |b|$, where $|a| \leq a_0$ and $|b| = |\Im(s)| \geq b_0$. So there exists a constant B_2 such that

$$\sum_{|\beta|=m} \|L_{-sr}^m(\chi_\beta)\|_\mu \leq B_2 |\Im(s)| \sum_{|\beta|=m} e^{-ar^m(z_\beta)}.$$

We now estimate the sum on the right. Naud estimates it directly, using a well known theorem [PP2, Bo]. However in [PP2] this theorem does not involve a sum over words $|\beta| = m$, but a sum over m -periodic points $f^m z = z$. In [Bo], it's not clear that the theorem holds in the way that Naud claims either, so we

show how to recover the estimate another way using the theorem as stated in [PP2]. Define a function

$$\varphi_m(a) = \left(\sum_{f^m z = z} e^{-ar^m(z)} \right)^{\frac{1}{m}}.$$

Theorem 3.9. ([PP2]) *Let $r : U \rightarrow \mathbb{R}$ be the first return map corresponding to the flow ϕ_t , let $f : U \rightarrow U$ be the expanding map corresponding to the flow and let $a \in I \subset \mathbb{R}$ where I is a closed, bounded set in \mathbb{R} containing h , the entropy of the flow. Let $P(-ar)$ be the pressure of $-ar$. Then we have $\varphi_m(a) \rightarrow e^{P(-ar)}$ uniformly in $a \in I$ as $m \rightarrow \infty$. Hence,*

$$\ln \varphi_m(a) = \frac{1}{m} \ln \sum_{f^m z = z} e^{-ar^m(z)} \rightarrow P(-ar) \text{ as } m \rightarrow \infty,$$

uniformly in $a \in I$.

Therefore for all $\varepsilon > 0$ there exists N such that for all $m \geq N$ and $a \in I$, we have $|\ln \varphi_m(a) - P(-ar)| < \varepsilon$. This gives us

$$\ln \varphi_m(a) \leq |\ln \varphi_m(a) - P(-ar)| + P(-ar) < \varepsilon + P(-ar),$$

which implies that $\varphi_m(a) \leq e^{\varepsilon + P(-ar)}$ for all $m \geq N$ and $a \in I$.

Theorem 3.10. *For all $\varepsilon > 0$ there exists $B_\varepsilon > 0$ such that for any real a in I , and for all $m \geq 1$ we have*

$$\left(\sum_{f^m z = z} e^{-ar^m(z)} \right) \leq B_\varepsilon e^{m(\varepsilon + P(-ar))}.$$

Proof. Let $b_\varepsilon = \max\{\varphi_m(a)^m : 0 < m < N, a \in A\}$, and let $B_\varepsilon = \max\{b_\varepsilon, 1\}$. Then the inequality holds. □

Recall that we can choose y_β , and hence $z_\beta \in U_\beta$ however we like. Since U_β may not have an m -periodic point, we cannot simply choose $f^m z_\beta = z_\beta$ and use the above theorem. However, U_β must have a periodic point of some higher order, because the periodic points of f are dense in U . We need a much larger constant to get the required inequality.

Definition 3.11. Let $p(\beta)$ be the smallest integer such that U_β has a $p(\beta)$ periodic point z_β i.e. $f^{p(\beta)} z_\beta = z_\beta$. Define $p(m)$ for any m to be the smallest integer such that for all words β of length m , there exists $z_\beta \in U_\beta$ with $f^{p(m)} z_\beta = z_\beta$. Equivalently, we define

$$p(m) = \text{lcm}\{p(\beta) : |\beta| = m\},$$

where lcm is the lowest common multiple of the set.

Now if we choose z_β so that $f^{p(m)}z_\beta = z_\beta$, then we have

$$\sum_{|\beta|=m} e^{-ar^{p(m)}(z_\beta)} = \sum_{f^{p(m)}z=z} e^{-ar^{p(m)}(z)}.$$

Finally we write $e^{-ar^m(z_\beta)}$ in terms of $e^{-ar^{p(m)}(z_\beta)}$ to get the estimate we need. For any $m \geq 1$, we have $p(m) \leq m + d$ for some integer constant d that depends only on the matrix A . This clearly follows from the irreducibility of A . We have

$$\begin{aligned} r^{p(m)}(z_\beta) &\leq r^{m+d}(z_\beta) = r(z_\beta) + \dots + r(f^{m+d-1}z_\beta) \\ &\leq r^m(z_\beta) + r(f^m z_\beta) + \dots + r(f^{m+d-1}z_\beta), \\ |r^{p(m)}(z_\beta) - r^m(z_\beta)| &\leq r^d(f^m z_\beta) \leq d|r|_\infty. \end{aligned}$$

So,

$$\begin{aligned} \sum_{|\beta|=m} e^{-ar^m(z_\beta)} &\leq \sum_{|\beta|=m} e^{-ar^{p(m)}(z_\beta)} |e^{-a(r^m(z_\beta) - r^{p(m)}(z_\beta))}| \\ &\leq \sum_{|\beta|=m} e^{-ar^{p(m)}(z_\beta)} e^{|a|d|r|_\infty} \\ &\leq e^{d|a||r|_\infty} \sum_{f^{p(m)}z=z} e^{-ar^{p(m)}(z_\beta)}. \end{aligned}$$

Now we can finally use Theorem 3.10. For all $\varepsilon > 0$, there exists $B_\varepsilon > 0$ such that

$$\sum_{|\beta|=m} e^{-ar^m(z_\beta)} \leq e^{d|a||r|_\infty} B_\varepsilon e^{(m+d)(\varepsilon + P(-ar))}.$$

We can group the constants together by defining $C_\varepsilon = C_5 B_2 B_\varepsilon e^{d|a||r|_\infty} e^{d(\varepsilon + P(-ar))}$. Then we get

$$\sum_{|\beta|=m} \|L_{-sr}^m(\chi_\beta)\|_\mu \leq C_5^{-1} C_\varepsilon |\Im(s)| e^{m(\varepsilon + P(-ar))}.$$

Combining this with the third particular estimate, and the inequality (3), we get the following:

For any set of points $x_1, \dots, x_k \in U_1, \dots, U_k$, and for all $\varepsilon > 0$, there exists a constant $C_\varepsilon > 0$ such that,

$$\left| Z_n(-sr) - \sum_{i=1}^k L_{-sr}^n \chi_i(x_i) \right| \leq C_\varepsilon |\Im(s)| \sum_{m=2}^n \left(\|L_{-sr}^{n-m}\|_\mu (\gamma^\mu)^m e^{m(\varepsilon + P(-ar))} \right).$$

and this proves Ruelle's Lemma.

4 Applying Ruelle's Lemma

Here we explain some of the context of Ruelle's Lemma, and show how it can be used in some cases to estimate the Ruelle zeta function. This was done by Pollicott and Sharp in [PS1, PS2] and later by Naud [Na] in different cases. Both estimates apply in less general cases than the conditions for Theorem 3.1. In particular they only apply when the stable and unstable manifolds are one-dimensional. Both estimates also depend on special cases of a very important result of Dolgopyat [Do1, Do2]. As an example of how Theorem 3.1 can be used, we show an estimate of the zeta function in the case considered by Naud [Na], which uses the following version of Dolgopyat's theorem.

Theorem 4.1. (*Dolgopyat*) *Let ϕ_t be a smooth transitive Anosov flow on a compact Riemannian manifold such that ϕ_t is non-integrable and has entropy h , and the local stable and unstable foliations are C^1 . Then there exists $C_6 > 0, \varepsilon_2 > 0, b_0 > 0$ and $0 < \rho < 1$ such that whenever $h - \varepsilon_2 \leq \Re(s) \leq h$ and $|\Im(s)| \geq b_0$,*

$$\|L_{-sr}^n\|_{\text{Lip}} \leq C_6 |\Im(s)| \rho^n, \quad \forall n \geq 0.$$

A similar estimate for open billiard flows was proved in [St1], and more general results were proved in [St2]. Since $W_\varepsilon^s(x)$ and $W_\varepsilon^u(x)$ are C^1 , the first return map is Lipschitz so we have $\mu = 1$.

Let $a = \Re(s) \in [h - \varepsilon_2, h]$ and let $|\Im(s)| \geq b_0$ as in the above theorem. By applying theorems 3.1 and 4.1, we get

$$\begin{aligned} |Z_n(s)| &\leq |Z_n(s) - \sum_{i=1}^k L_{-sr}^n(\chi_i)(x_i)| + |\sum_{i=1}^k L_{-sr}^n(\chi_i)(x_i)| \\ &\leq C_\varepsilon |\Im(s)| \sum_{m=2}^n \left(\|L_{-sr}^{n-m}\|_{\text{Lip}} \gamma^m e^{m(\varepsilon + P(-ar))} \right) + \sum_{i=1}^k \|L_{-sr}^n\|_{\text{Lip}} \\ &\leq C_\varepsilon |\Im(s)| \sum_{m=2}^n \left(C_6 |\Im(s)| \rho^{n-m} \gamma^m e^{m(\varepsilon + P(-ar))} \right) + k C_6 |\Im(s)| \rho^n \\ &\leq C_6 C_\varepsilon |\Im(s)|^2 \rho^n \sum_{m=2}^n \left(\frac{\gamma}{\rho} e^{\varepsilon + P(-ar)} \right)^m + k C_6 |\Im(s)| \rho^n. \end{aligned}$$

Although the ρ in Theorem 4.1 is fixed, the inequality in the theorem still holds for larger ρ , as long as $\rho < 1$. Thus we can choose ρ so that $1 > \rho > \gamma$. Note that $P(-hr) = 0$, so by taking ε small enough, $a = \Re(s)$ sufficiently close to h , say $|a - h| < \delta$ for some $\delta > 0$, and $\frac{\gamma}{\rho}$ sufficiently small, we can have

$$\frac{\gamma}{\rho} e^{\varepsilon + P(-ar)} < 1.$$

This simplifies the inequality to

$$|Z_n(s)| \leq C_6 C_\varepsilon |\Im(s)|^2 \rho^n + k C_6 |\Im(s)| \rho^n \leq C_7 |\Im(s)|^2 \rho^n,$$

where $C_7 = C_6 C_\varepsilon + k C_6 b_0^{-1}$ is simply another constant. Now we can obtain an estimate for $\zeta(s)$. The Ruelle zeta function [PS1] is given by

$$\zeta(s) = \exp \left(\sum_{n=1}^{\infty} \frac{1}{n} Z_n(-sr) \right).$$

So we can continue this zeta function analytically to the domain

$$\{s : |\Re(s) - h| < \min\{\varepsilon_2, \delta\} \text{ and } |\Im(s)| \geq b_0\},$$

for some $\varepsilon_2, b_0 > 0$. In this region we have

$$\begin{aligned} \exp \left(- \sum_{n=1}^{\infty} \frac{1}{n} |Z_n(-sr)| \right) &\leq |\zeta(s)| \leq \exp \left(\sum_{n=1}^{\infty} \frac{1}{n} |Z_n(-sr)| \right) \\ \exp \left(-C_7 |\Im(s)|^2 \sum_{n=1}^{\infty} \frac{1}{n} \rho^n \right) &\leq |\zeta(s)| \leq \exp \left(C_7 |\Im(s)|^2 \sum_{n=1}^{\infty} \frac{1}{n} \rho^n \right) \\ \exp \left(-C_7 |\Im(s)|^2 \ln \left(\frac{1}{1-\rho} \right) \right) &\leq |\zeta(s)| \leq \exp \left(C_7 |\Im(s)|^2 \ln \left(\frac{1}{1-\rho} \right) \right). \end{aligned}$$

This demonstrates the application of Ruelle's Lemma to estimating the zeta function.

References

- [Bo] R. Bowen, Equilibrium states and the ergodic theory of Anosov diffeomorphisms, Lecture Notes in Mathematics (1975), 470.
- [Do1] D. Dolgopyat, On statistical properties of geodesic flows on negatively curved surfaces, PhD Thesis, Princeton University, 1997.
- [Do2] D. Dolgopyat, On decay of correlations in Anosov flows, *Ann. of Math.* **147** (1998), 357-390.
- [Ha] B. Hasselblatt, Regularity of the Anosov splitting and of horospheric foliations. *Ergod. Th. & Dynam. Sys.* **14** (1994), 645-666.
- [Na] F. Naud, Expanding Maps on Cantor sets and analytic continuation of zeta functions. *Ann. Sci. École Norm. Sup.* **38** (2005), 116-153.
- [PP1] W. Parry and M. Pollicott, Zeta Functions and the Periodic Orbit Structure of Hyperbolic Dynamics, *Astérisque* (1990).
- [PP2] W. Parry and M. Pollicott, Zeta Functions and the periodic orbit structure of hyperbolic dynamics. Page 74.
- [PS1] M. Pollicott and R. Sharp, Exponential Error Terms for Growth Functions on Negatively Curved Surfaces. *Amer. J. Math.* **120** (1998), 1019-1042.
- [PS2] M. Pollicott and R. Sharp, Error Terms for Closed Orbits of Hyperbolic Flows. *Ergodic Theory Dynam. Systems*, **21**(2):545-562, 2001.

- [Ru] D. Ruelle, An Extension of the Theory of Fredholm Determinants, Publ. Math. (IHES). **72** (1990), 175-193
- [St1] L. Stoyanov, Spectrum of the Ruelle operator and exponential decay of correlations for open billiard flows, Amer. J. Math. **123** (2001), 715-759
- [St2] L. Stoyanov, Spectra of Ruelle transfer operators for Axiom A flows, Preprint (arXiv: math.DS:0810.1126).
- [W] P. Walters, An introduction to ergodic theory, Springer, New York, 1982.
- [WW] J.H. Wells and L.R. Williams, Embeddings and Extensions in Analysis, Berlin, Springer, 1975.

UNIVERSITY OF WESTERN AUSTRALIA, CRAWLEY WA, 6009, AUSTRALIA
E-mail address: *paul.e.wright@uwa.edu.au*