

Law of the Iterated Logarithm for the random walk on the infinite percolation cluster

Hugo Duminil-Copin

September 2008

ABSTRACT : We show that random walks on the infinite supercritical percolation clusters in \mathbb{Z}^d satisfy the usual Law of the Iterated Logarithm. The proof combines Barlow's Gaussian heat kernel estimates and the ergodicity of the random walk on the environment viewed from the random walker as derived by Berger and Biskup.

1 Introduction

Asymptotic properties of random walks in \mathbb{Z}^d are very well-understood. Their convergence to d -dimensional Brownian motions and their almost sure behavior (such as the law of the iterated logarithm) have been derived decades ago. A natural question to ask is what happens to random walks on graphs that are in some sense perturbations of \mathbb{Z}^d . One of the first examples to consider is to look at the random graph obtained by taking the infinite cluster $\mathcal{C} = \mathcal{C}(\omega)$ of a supercritical percolation process. One “perturbs” the original lattice by removing some edges independently. Various large-scale properties of this infinite graph have been studied with techniques such as coarse-graining. One of the most natural questions is to look at random walk on this cluster and to study its behavior.

One can for instance consider the continuous-time simple random walk (CTSRW) on \mathcal{C} . This is the process X^ω that waits an exponential time of mean 1 at each vertex x and jumps along one of the open edges e adjacent to x , with each edge chosen with equal probability. This process has been studied in a number of papers. Grimmett, Kesten, and Zhang ([5],1993) proved that X^ω is almost surely recurrent if $d = 2$ and transient if $d \geq 3$. Barlow ([2],2004) proved Gaussian estimates for X^ω . An invariance principle in every dimension has been proved independently by Berger and Biskup in ([3],2004) and by Mathieu and Piatnitski ([8],2004). Before that, Sidoravicius and Sznitman proved this result for $d \geq 4$ ([9], 2004). All these results show that a property that holds for random walk on \mathbb{Z}^d still holds for random walk on the infinite supercritical percolation cluster.

It is natural to ask if this is still valid if one looks for instance at almost sure properties of the random walk (recall that almost sure properties often describe the behavior of the walk at exceptional times). Our goal in the present note is to show that it is indeed the case for the law of the iterated logarithm (LIL).

Theorem 1.1 Consider $d \geq 2$ and suppose that $p > p_c$, where $p_c = p_c(d)$ is the critical bond percolation probability in \mathbb{Z}^d . Then, there exists a positive and finite constant $c(p, d)$, such that for almost all realization of percolation with parameter p , for all x in the infinite cluster \mathcal{C} , the continuous-time random walk X^w started from x satisfies almost surely the following LIL :

$$\limsup_{t \rightarrow \infty} \frac{|X_t^\omega|}{\sqrt{t \log \log t}} = c(p, d).$$

Here and throughout the paper, $|x| = |x|_1 = \sum_{j=1}^d |x_j|$ stands for the L^1 norm of $x = (x_1, \dots, x_d) \in \mathbb{Z}^d$. Our proof can trivially be adapted to other norms (this would just change the value of the constant). Note also that we are studying almost sure properties of the walk, so that the annealed and quenched statements are identical here (once we say that the constant $c(p, d)$ does not depend on the environment).

The main ingredients of our proof are the Gaussian bounds derived in Barlow [2] and the ergodicity of Kipnis-Varadhan's [7] random walk on the environment as seen from the random walker derived by Berger and Biskup in [3]. These two results have in fact been instrumental in the (much more difficult) derivation of the invariance principle for this random walk.

The paper is organized as follows. In Section 2, we will show that one can find positive and finite $c_1(p, d)$ and $c_2(p, d)$ such that almost surely

$$c_1(p, d) \leq \limsup_{t \rightarrow \infty} \frac{|X_t^\omega|}{\sqrt{t \log \log t}} \leq c_2(p, d).$$

This will be based on the Gaussian estimates derived in [2]. The upper bound is an easy application of the Borel-Cantelli Lemma whereas the proof of the lower bound will use the Markov property and the fact that one can apply Gaussian bounds uniformly for x in a ball of sufficiently large radius depending on t .

In Section 3, we derive a Zero-one Law for the limit of a discrete analog of the CTSRW. The main ingredient will be the ergodicity of a certain shift T , related to Kipnis-Varadhan's random walk on the environment. It has been proved by Noam Berger and Marek Biskup [3] that this shift T is ergodic. Translating properties of the random walk in terms of this shift will allow us to derive a Zero-one law for the limsup in the LIL for this discrete-time random walk.

Finally, in Section 4, we conclude by checking that the time-scales of the discrete-time random walk and of the continuous-time random walk are comparable.

2 Weak LIL for the continuous random walk

We will consider Bernoulli bond percolation of parameter p on \mathbb{Z}^d defined on a probability space $(\Omega, \mathcal{F}, \mathbb{P}_p)$. It is well known (Grimmett [6]) that there exists $p_c \in (0, 1)$ such that when $p > p_c$ there is a unique infinite open cluster, that we denote by \mathcal{C} . For \mathbb{P}_p almost every environment $\omega \in \Omega$ and $x \in \mathcal{C}$, we define a CTSRW $X^\omega = (X_t^\omega, t \geq 0)$ started from x under the probability measure \mathbb{P}_ω^x . **In the whole paper, we fix $p > p_c$ and $d \geq 2$.**

Because of translation-invariance of our problem (and because we are dealing with almost sure properties), we can restrict ourselves to the case $x = 0$ and work with the probability measure $\tilde{\mathbb{P}}_p = \mathbb{P}_p(\cdot | 0 \in \mathcal{C})$ and $X_0^\omega = 0$. We will use the notation $\Phi(t) = \sqrt{t \log \log t}$ for all $t > e$.

We now recall Barlow's Gaussian estimates. The first one uses the chemical distance d_ω (or graph distance) on \mathcal{C} . For every x and y in \mathcal{C} , $d_\omega(x, y)$ is the length of the shortest path between x and y that uses only edges in \mathcal{C} . For every integer n and $x \in \mathcal{C}$, $\mathcal{B}_\omega(x, n)$ will denote the ball of radius n and of center x for distance d_ω .

Proposition 2.1 (Barlow, [2]) *There exist two constants $a_1 = a_1(p, d)$ and $a_2 = a_2(p, d)$ such that for every $\gamma > 0$, there exists a finite random variable M_γ satisfying for almost every environment ω :*

$$\text{for all } n \geq M_\gamma(\omega), \quad \mathbb{P}_\omega^0(\max_{k \in [0, n]} d_\omega(0, X_k^\omega) > \gamma \Phi(n)) \leq a_1 \exp\left(-a_2 \frac{(\gamma \Phi(n))^2}{n}\right).$$

Recall that this statement holds for a general class of graphs (see Proposition 3.7 of Barlow [2]); percolation estimates (see Theorem 2.18 and Lemma 2.19 of Barlow [2]) show that the percolation cluster belongs to this class. The other result that we will use is the Gaussian bound itself :

Theorem 2.2 (Barlow, [2]) *There exist finite constants c_1, \dots, c_8 and $\epsilon > 0$ only depending on p and d that satisfy the following property. There exists a random variable S_0 with $\tilde{\mathbb{P}}_p(S_0 \geq n) \leq c_7 \exp(-c_8 n^\epsilon)$ and for almost every environment ω such that $0, y \in \mathcal{C}, t \geq 1$:*

(1) *The transition density $p_t^\omega(0, y)$ of X^ω satisfies the Gaussian bound*

$$c_1 t^{-d/2} e^{-c_2 |y|^2/t} \leq p_t^\omega(0, y) \leq c_3 t^{-d/2} e^{-c_4 |y|^2/t} \text{ for } t \geq S_0(\omega) \vee |y|.$$

(2) *$c_5 n^d \leq \text{Vol}(\mathcal{B}_\omega(0, n)) \leq c_6 n^d$ for $n \geq S_0(\omega)$.*

Note that translation invariance makes possible for each $x \in \mathbb{Z}^d$, a random variable S_x satisfying the analogous conditions (with the same constants $c_1, \dots, c_8, \epsilon$) where one just replaces the origin 0 by x (and therefore replaces y by $x + y$).

Let remark that there is no uniform Gaussian bounds for every $x, y \in \mathcal{C}$ and every $t > 0$ because (almost surely) every finite graph is actually embedded somewhere in the infinite cluster. We can now derive almost sure upper and lower bounds for our limsup.

Proposition 2.3 (Upper bound) *There exists a finite $c_+ = c_+(p, d)$ such that for almost every environment ω ,*

$$\mathbb{P}_\omega^0 \text{ a.s. } \limsup_{t \rightarrow \infty} \frac{|X_t^\omega|}{\Phi(t)} \leq c_+.$$

Proof : Fix ω an environment containing 0. The proof goes along the same lines as in the Brownian case. Let $\gamma > 0$, and define the following events :

$$A_n^\omega = \left\{ \max_{k \in [0, 2^n]} d_\omega(0, X_k^\omega) > \gamma \Phi(2^n) \right\}.$$

Proposition 2.1 shows that for all n large enough,

$$\mathbb{P}_\omega^0(A_n^\omega) \leq a_1 \exp\left(-a_2 \frac{(\gamma\Phi(2^n))^2}{2^n}\right) \leq 2a_1 n^{-a_2\gamma^2}.$$

Providing γ large enough, the Borel-Cantelli Lemma claims that almost surely A_n^ω holds finitely often. Using the fact that $|\cdot| \leq d_\omega(0, \cdot)$, we get that for n large enough, $\max_{k \in [0, 2^n]} |X_k^\omega| < \gamma\Phi(2^n)$. We conclude that for n large enough, $|X_n^\omega| < 2\gamma\Phi(n)$. \square

Proposition 2.4 (Lower bound) *There exists a positive $c_- = c_-(p, d)$ such that for almost every environment ω ,*

$$\mathbb{P}_\omega^0 \text{ a.s.}, c_- \leq \limsup_{t \rightarrow \infty} \frac{|X_t^\omega|}{\Phi(t)}.$$

Let first present the outline of the proof. Consider $q > 1$ and $\gamma > 0$ (we will choose their values later). As in the Brownian case, set $D_n^\omega = X_{q^n}^\omega - X_{q^{n-1}}^\omega$. We have $|X_{q^n}^\omega| \geq |D_n^\omega| - |X_{q^{n-1}}^\omega|$. Using the upper bound, we obtain that almost surely, for n large enough :

$$|X_{q^n}^\omega| \geq |D_n^\omega| - 2c_+\Phi(q^{n-1}). \quad (2.1)$$

Because $\Phi(q^{n-1}) \leq q^{-1/2}\Phi(q^n)$, the second term can be chosen much smaller than $\Phi(q^n)$, providing q large enough. Then, in order to prove the result, it is enough to bound D_n^ω from below. Define the events $C_n^\omega = \{|D_n^\omega| > \gamma\Phi(q^n)\}$. If these events hold for infinity many n almost surely, then we are done. We define the σ -fields $\mathcal{F}_n^\omega = \sigma(X_k^\omega, k \leq q^n)$. We will apply the Borel-Cantelli Lemma generalized to dependent events (see Durrett [4], chapter 4, paragraph 4.3). We therefore need to prove that

$$\mathbb{P}_\omega^0 \text{ a.s.} \sum_{n \geq 1} \mathbb{E}_\omega^0[C_n^\omega | \mathcal{F}_{n-1}^\omega] = \infty.$$

Using the Markov property and Gaussian bounds, we will be able to find a lower bound for $\mathbb{E}_\omega^0[C_n^\omega | \mathcal{F}_{n-1}^\omega]$. In order to apply these bounds, we need to control not only S_0 (from Theorem 2.2) but also S_x for $x = X_{q^{n-1}}^\omega$. We first prove that it is indeed possible, using Gaussian estimates and the upper bound.

Lemma 2.5 *Let $\gamma > 0$, for almost every environment ω we have almost surely $S_{X_n^\omega} \leq \gamma\Phi(n)$ for n large enough.*

Proof : Let $\gamma > 1$. Define for each integer n the set

$$B_n = \{\exists y \in B(0, 2c_+\Phi(n)) \text{ s.t. } S_y \geq \gamma\Phi(n)\}.$$

where S_y is the random variable of Theorem 2.2. The Theorem yields

$$\tilde{\mathbb{P}}_p(B_n) \leq \text{Vol}(B(0, 2c_+\Phi(n))) d_1 \exp(-d_2(\gamma\Phi(n))^\epsilon).$$

The right-hand side of the inequality is summable, so that (by Borel-Cantelli) B_n holds for a finite number of n for almost every environment. But almost surely, X_n^ω is less than $2c_+\Phi(n)$ for n large enough. Combining these two facts, we obtain the claim.

Proof of Proposition 2.4 : Let $q, \gamma > 0$ and $\kappa > 0$ such that $c_5 \kappa^d > c_6 + 1$. Note that κ does not depend on γ and q . Set $t_n = q^n - q^{n-1}$. By the Markov property, we get for $n \geq 1$,

$$\mathbb{E}_\omega^0(C_n^\omega | \mathcal{F}_{n-1}^\omega) = \mathbb{P}_{\omega_n}^0[\gamma\Phi(q^n) < X_{t_n}^{\omega_n}] \geq \mathbb{P}_{\omega_n}^0[\gamma\Phi(q^n) < X_{t_n}^{\omega_n} < \kappa\gamma\Phi(q^n)] = G_n(\omega_n)$$

where $\omega_n = \tau_{X_{q^{n-1}}^\omega}(\omega)$ (τ_x is the shift defined by $(\tau_x\omega)_y = \omega_{x+y}$) and :

$$G_n(\omega) = \mathbb{P}_\omega^0[\gamma\Phi(q^n) < X_{t_n}^\omega < \kappa\gamma\Phi(q^n)].$$

The function G_n is well-defined and measurable. If $\mathcal{A}_n(\omega)$ is the annulus

$$\mathcal{A}_n(\omega) = \{z \in \mathcal{C}, \text{ s.t. } \gamma\Phi(q^n) < |z| < \kappa\gamma\Phi(q^n)\},$$

we find by definition of the transition density $G_n(\omega) = \sum_{z \in \mathcal{A}_n(\omega)} p_{t_n}^\omega(0, z)$. We deduce :

$$\mathbb{E}_\omega^0(C_n^\omega | \mathcal{F}_{n-1}^\omega) = \sum_{z \in \mathcal{A}_n(\omega_n)} p_{t_n}^{\omega_n}(0, z). \quad (2.2)$$

Using Lemma 2.5, we know that almost surely there exists N large enough such that for every n larger than N , $S_0(\omega_n) = S_{X_{q^{n-1}}^\omega}(\omega) \leq \gamma\Phi(q^{n-1}) \leq t_n$. For $n \geq N$, one can use Gaussian estimates of Theorem 2.2 for every $z \in \mathcal{A}_n(\omega_n)$, we get for such a z :

$$p_{t_n}^{\omega_n}(0, z) \geq c_1 t_n^{-d/2} \exp\left(-\frac{c_2(\kappa\gamma\Phi(q^n))^2}{t_n}\right) \geq c_1 t_n^{-d/2} n^{-c_2(\kappa\gamma)^2} \quad (2.3)$$

Using again the same Lemma, Theorem 2.2 yields that the volume growth property holds for $S_n(\omega_n)$. Recalling the definition of κ , we find :

$$\text{Vol}(\mathcal{A}_n(\omega_n)) \geq (\gamma\Phi(q^n))^d \geq \gamma^d t_n^{d/2} \quad (2.4)$$

Combining (2.3) and (2.4) in (2.2), we obtain that there exists a constant $c > 0$ such that almost surely for n large enough :

$$G_n(\omega_n) \geq cn^{-c_2(\kappa\gamma)^2}$$

Providing γ small enough, we can use the generalized Borel-Cantelli Lemma (e.g. [4]). We get that almost surely, there exist infinitely many integers n such that $|D_n^\omega| > \gamma\Phi(q^n)$. If $q > 0$ is taken large enough (κ, γ and c_2 are not depending on q), we can use the inequality (2.1) to prove that almost surely :

$$|X_{q^n}^\omega| \geq \gamma\Phi(q^n) - 2c_+ q^{-1/2} \Phi(q^n) > \frac{\gamma}{2} \Phi(q^n)$$

for infinitely many n , which is the claim.

Remark 1 : In order to bound the sum in (2.2) from below, Gaussian bounds were not sufficient. Without the volume growth property, the annulus could contain only few elements. Even if the exponential term is not too small (typically of order n^{-s} for s small), the term $t_n^{-d/2}$ (which corresponds to $t^{-d/2}$ for the Brownian motion) could be very small and make the series become summable. The cardinality of the annulus was critical in order to balance out this term.

Remark 2 : our goal was to obtain a result in L^1 norm. Unfortunately, the natural distance on graphs is the chemical distance $d_\omega(\cdot, \cdot)$. In the bound from below, this does not create any trouble because of the trivial inequality $|x| \leq d_\omega(0, x)$. But it could happen that the chemical distance is much bigger than the L^1 norm. The proof of Theorem 2.18 in Barlow [2] precisely deals with this issue thanks to a result by Antal and Pisztora [1] that shows that the chemical distance on \mathcal{C} and the L^1 norm are not that different on a supercritical percolation cluster.

3 Zero-one Law for the blind random walk

In the present section, we will consider discrete time random walks. We first introduce the two random walks we will use. Then we recall an ergodicity result proved in [3] and we derive the Zero-one Law. Our proofs are rather direct applications of the ergodicity statement of [3].

For each $x \in \mathbb{Z}^d$, let τ_x be the shift from Ω in Ω defined by $(\tau_x \omega)_y = \omega_{y+x}$. For each ω , let Y_n^ω be the simple random walk (called **blind random walk**) on \mathcal{C} started at the origin. At each unit of time, the walk picks a neighbor at random and if the corresponding edge is occupied, the walk moves to this neighbor. Otherwise, it does not move. This random walk may seem less natural than the random walk that chooses randomly one of the accessible neighbors and jumps to it, but this blind random walk preserves the uniform measure on \mathcal{C} , so that the stationary measure on the environment as seen from the walker turns out to be simpler.

It is well known (cf Kipnis and Varadhan [7]) that the Markov chain $(Y_n^\omega)_{n \geq 0}$ induces a Markov chain on Ω (the so-called **Markov chain on the environment**), that can be interpreted as the trajectory of "environment viewed from the perspective of the walk". It is defined as

$$\omega_n(\cdot) = \omega(\cdot + Y_n^\omega) = \tau_{Y_n^\omega} \omega(\cdot).$$

One can describe the chain (ω_n) as follows. At each step n , one chooses one of the $2d$ neighbors of the origin at random and calls it e . If the corresponding edge is closed for ω_n , then $\omega_{n+1} = \omega_n$, otherwise $\omega_{n+1}(\cdot) = \tau_e \circ \omega_n$, where $\tau_e \circ \omega(\cdot) = \omega(\cdot - e)$.

It is straightforward to check that the probability measure $\tilde{\mathbb{P}}_p$ is a reversible and therefore stationary measure for the Markov chain (ω_n) . This allows us to extend our probability space to $\Xi = \Omega^{\mathbb{Z}}$ (endowed with the product σ -algebra $\mathcal{H} = \mathcal{F}^{\otimes \mathbb{Z}}$) and to

define ω_n also for negative n 's in such a way that that the family $(\omega_n, n \in \mathbb{Z})$ is stationary. Let μ denote the probability measure associated to the Markov chain.

Note that under the measure μ , and for all $n \in \mathbb{Z}$, the law of $(\omega_n, \omega_{n+1}, \dots)$ is identical to that of $(\omega_0, \omega_1, \dots)$. On the other hand, the marginal law of ω_0 (still under μ) is $\tilde{\mathbb{P}}_p$. One then defines $T : \Xi \rightarrow \Xi, \bar{\omega} \mapsto T\bar{\omega}$ to be $(T\bar{\omega})_n = \bar{\omega}_{n+1}$.

Theorem 3.1 (Berger, Biskup, [3]) *T is ergodic with respect to μ . In other words, for all $A \in \mathcal{H}$, if $T^{-1}(A) = A$, then $\mu(A)$ is equal to 0 or 1.*

We refer to the paper of Berger and Biskup [3] for proofs. Define for every $a > 0$ and ω the event :

$$A_\omega(a) = \left\{ \limsup_{n \rightarrow \infty} \frac{|Y_n^\omega|}{\Phi(n)} > a \right\}.$$

Let now state and prove a consequence of this ergodicity for our law of the iterated logarithm :

Corollary 3.2 (Zero-one Law) *Let $a \geq 0$. The probability that*

$$B_a = \{\mathbb{P}_\omega^x \text{ a.s. } A_\omega(a) \text{ holds for all } x \in \mathcal{C}\}$$

is equal to 0 or to 1.

Proof : Our goal is to use the ergodicity of the environment and to note that the considered event corresponds to a T -invariant set in Ξ . Let $a \geq 0$ and define the function F on Ω by :

$$F(\omega) = \mathbb{P}_\omega^0(A_\omega(a))$$

This function is well-defined and measurable. Let fix the environment ω for a little while and denote $\omega_n = \tau_{Y_n^\omega} \omega$. We claim that $(F(\omega_n))_n$ is a martingale with respect to the filtration \mathcal{F}_n associated to the process Y_n^ω . Indeed, the Markov property yields

$$F(\omega_n) = \mathbb{P}_\omega^0(A_\omega(a) | \mathcal{F}_n).$$

This martingale is bounded and therefore converges almost surely as $n \rightarrow \infty$. Moreover, it converges to the indicator function of $A_\omega(a)$ because this event is clearly in $\mathcal{F}_\infty = \sigma(\cup_{n \geq 0} \mathcal{F}_n)$.

By taking the Cesaro mean and then integrating it with respect to ω (and using the fact that the probabilities are bounded by 1), we get that

$$\tilde{\mathbb{E}}_0 \left[\mathbb{P}_\omega^0 \left(\left| \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=0}^{N-1} F(\omega_n) - 1_{A_\omega(a)} \right| \right) \right] = 0$$

On the other hand, F can be viewed as a measurable function on Ξ . The ergodicity of μ implies that for μ almost every $\bar{\omega}$:

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N F(\bar{\omega}_n) = \int F d\mu$$

Let recall that $\bar{\omega}$ has same law under μ as $(\omega_n)_n$ under $\tilde{\mathbb{E}}_0[\mathbb{P}_\omega^0(\cdot)]$. We deduce that the limit $1_{A_\omega(a)}$ is (up to a set of zero measure) constant. Since it is an indicator function, this means that either the corresponding event is almost surely true, or almost surely wrong.

4 The Law of the Iterated Logarithm

We can now derive the Law of the Iterated Logarithm. Let first note that the previous corollary immediately implies that for a fixed $p > p_c$, there exists a constant $c'(p, d) \in [0, \infty]$ such that for almost every environment ω , the blind random walk satisfies

$$\limsup_{n \rightarrow \infty} \frac{|Y_n^\omega|_1}{\Phi(n)} = c'(p, d)$$

almost surely (just choose $c'(p, d)$ to be the supremum of the set of a 's such that the event B_a is almost surely satisfied).

Our next goal is to show that the time scales for the two random walks are comparable. Let $\omega \in \Omega_0$, define the real random variable $(T_n^\omega)_n$ by $T_0^\omega = 0$ and

$$T_{n+1}^\omega = \inf \{t > T_n^\omega, X_t^\omega \neq X_{T_n^\omega}^\omega\}.$$

Clearly, the Law of Large Numbers implies that for all $\omega \in \Omega_0$, $T_{n+1}^\omega \sim n$ almost surely.

Let $\omega \in \Omega_0$, define in the same way the random variable (U_n^ω) by $U_0^\omega = 0$ and

$$U_{n+1}^\omega = \inf \{p > U_n^\omega, Y_p^\omega \neq Y_{U_n^\omega}^\omega\}.$$

The $(U_{n+1}^\omega - U_n^\omega)_n$ are not i.i.d. anymore. Conditionally on the environment and on the past up to the n -th jump of Y^ω , the law of $U_{n+1}^\omega - U_n^\omega$ is geometric and depends on the number $I(n)$ of incoming open edges at $Y^\omega U_n^\omega$ (its mean is some function $f(I(n))$).

Ergodicity ensures that almost surely and for each $k \leq 2d$,

$$\frac{1}{n} \sum_{j=1}^n 1_{I(j)=k} \rightarrow i(k)$$

where $i(k)$ denotes the μ -probability that ω_0 has k incoming open edges at the origin.

Using the Law of Large Numbers for sums of independent geometric random variables of mean $f(k)$ for each k , we get readily that for almost all $\omega \in \Omega_0$,

$$\mathbb{P}_\omega^0 \text{ a.s. } U_{n+1}^\omega/n \rightarrow \sum_{k=1}^{2d} i(k)f(k) = \alpha_p^{-1}.$$

This last quantity is clearly positive and finite.

We can now conclude the proof of the Law of the Iterated Logarithm for the continuous time random walk.

Proof of Theorem 1.1 : Consider the natural coupling for which X_t^ω and Y_n^ω have the same trajectories. More precisely, if we consider the **myopic random walk** $(Z_n^\omega)_n$ that jumps at each time, choosing uniformly a neighbor, defined on a probability space $(\Omega_\omega, \mathcal{F}_\omega, \mathbb{P}_\omega^0)$. Assume there exists an independent family $(T_i)_{i \in \mathbb{Z}_+}$ of iid exponential mean time 1 random variables and $(S_x^\omega)_{x \in \mathbb{Z}^d}$ an independent family of independent random variables such that S_x^ω is a geometrical of parameter $n_x^\omega/(2d)$ where n_x^ω is the number of adjacent open edges of x for the configuration ω .

Define $T_p^\omega = \sum_{k=0}^{p-1} T_i$ and $n^\omega(t) = \sup \{p, T_p^\omega \leq t\}$. Then we can write the continuous time random walk as follows

$$X_t^\omega = Z_{n^\omega(t)}^\omega \quad \forall t \geq 0.$$

Now, consider $U_p^\omega = \sum_{k=0}^{p-1} S_{Z_k^\omega}^\omega$ and $m^\omega(n) = \sup \{p, U_p^\omega \leq n\}$. Then we can write the blind random walk as follow :

$$Y_n^\omega = Z_{m^\omega(n)}^\omega \quad \forall n \geq 0.$$

Because of the estimates of the time-scales of our two walks, we get that

$$\limsup_{t \rightarrow \infty} \frac{|X_t^\omega|}{\Phi(t)} = \limsup_{t \rightarrow \infty} \frac{|Z_{n^\omega(t)}^\omega|}{\Phi(n^\omega(t))} = \limsup_{n \rightarrow \infty} \frac{|Z_n^\omega|}{\Phi(n)}$$

and that

$$\limsup_{n \rightarrow \infty} \frac{|Y_n^\omega|}{\Phi(n)} = \limsup_{n \rightarrow \infty} \frac{|Z_{m^\omega(n)}^\omega|}{\Phi(\alpha_p m^\omega(n))} = \frac{1}{\sqrt{\alpha_p}} \limsup_{n \rightarrow \infty} \frac{|Z_n^\omega|}{\Phi(n)}.$$

From these two equalities, we deduce that

$$\limsup_{t \rightarrow \infty} \frac{|X_t^\omega|}{\Phi(t)} = \frac{1}{\sqrt{\alpha_p}} \limsup_{n \rightarrow \infty} \frac{|Y_n^\omega|}{\Phi(n)} \text{ a.s..}$$

The theorem follows readily. □

Note that this also show that the Law of the Iterated Logarithm holds for the blind and the myopic random walks.

Acknowledgements : This paper was written during my stay at the University of British Columbia, I would like to thank M.T. Barlow, who first taught me about this question, for his availability and the advice he gave me during my whole stay. I would also like to thank W. Werner for his careful reading of this paper and his numerous suggestions.

Références

- [1] P. Antal and A. Pisztor. On the chemical distance for supercritical bernoulli percolation. *Ann. Probab.*, 2(24), 1996.
- [2] M.T. Barlow. Random walks on supercritical percolation clusters. *Ann. Probab.*, 4(32), 2004.
- [3] N. Berger and M. Biskup. Quenched invariant principles for simple random walk on percolation clusters. *Probab. Theory Rel. Fields*, (137), 2007.
- [4] R. Durrett. *Probability : Theory and Examples, second edition*. Duxbury Press, 1989.

- [5] H. Kesten G. Grimmett and Y. Zhang. Random walk on the infinite cluster of the percolation model. *Probab. Theory Rel. Fields*, 1(96), 1993.
- [6] G.R. Grimmett. *Percolation (Second edition)*. Springer-Verlag, 1999.
- [7] C. Kipnis and S.R.S. Varadhan. A central limit theorem for additive functionals of reversible markov processes and applications to simple exclusions. *Commun. Math. Phys.*, 1(104), 1986.
- [8] P. Mathieu and A.L. Piatnitski. Quenched invariant principles for random walks on percolation clusters. *Proc. R. Soc. A*, (463), 2007.
- [9] V. Sidoravicius and A.S. Sznitman. Quenched invariant principles for walks on clusters of percolation or among random conductances. *Probab. Theory Rel. Fields*, 2(129), 2004.

DEPARTMENT OF MATHEMATICS
UNIVERSITY OF BRITISH COLUMBIA
VANCOUVER, BRITISH COLUMBIA, CANADA

DMA, ECOLE NORMALE SUPÉRIEURE
45 RUE D'ULM, 75230 PARIS CEDEX 05, FRANCE
E-MAIL : hugo.duminil@ens.fr