

Topology of Conic Bundles - II

Nitin Nitsure

24 February 1996

Tata Institute of Fundamental Research, Mumbai 400 005, India.

e-mail: nitsure@tifrvax.tifr.res.in

1991 Mathematics subject classification : 14F20, 13A20, 12G05

Abstract

For conic bundles on a smooth variety (over a field of characteristic $\neq 2$) which degenerate into pairs of distinct lines over geometric points of a smooth divisor, we prove a theorem which relates the Brauer class of the non-degenerate conic on the complement of the divisor to the covering class (Kummer class) of the 2-sheeted cover of the divisor defined by the degenerate conic, via the Gysin homomorphism in étale cohomology. This theorem is the algebro-geometric analogue of a topological result proved earlier.

1 Introduction

Let X be a smooth scheme over a field F of characteristic $\neq 2$, and let $C \rightarrow X$ be a conic bundle on X , whose discriminant defines a smooth divisor $Y \subset X$ with multiplicity τ (which, if Y is not irreducible, will consist of a positive integer τ_i for each component Y_i of Y). Let $\beta \in H^2(X, \mu_2)$ be the Brauer class of the restriction of C to $X - Y$, which is a P^1 fibration which is étale locally trivial (all cohomologies are with respect to étale topology unless otherwise indicated). Suppose that over each geometric point of Y , the fiber of C consists of two distinct projective lines meeting at a point. Therefore, the relative Hilbert scheme of lines in $C|_Y \rightarrow Y$ is a two sheeted finite étale cover $\tilde{Y} \rightarrow Y$. Let $\alpha \in H^1(Y, \mu_2)$ be its covering class ('Kummer class'). By smoothness, we have a Gysin homomorphism $H^2(X - Y, \mu_2) \rightarrow H^1(Y, \mu_2)$. We prove here the following

Theorem 1.1 *Under the Gysin homomorphism $H^2(X - Y, \mu_2) \rightarrow H^1(Y, \mu_2)$, the Brauer class β of the P^1 fibration on $X - Y$ maps to $\tau\alpha$, where τ is the vanishing multiplicity of the discriminant and α is the cohomology class of the two sheeted finite étale cover $\tilde{Y} \rightarrow Y$.*

A topological version of this result was proved in [Ni] for topological conic bundles on manifolds, which is equivalent for complex algebraic conic bundles to the above

result. This is because there is a natural isomorphism between étale cohomology with finite constant coefficients (in this case, coefficients $\mathbb{Z}/(2)$) and the corresponding singular cohomology, which commutes with the two Gysin.

The theorem is proved below in two steps. In section 2, we reduce it to proving a purely algebraic lemma (Lemma 2.1 below, which we call as the ‘main lemma’) over a discrete valuation ring. In section 3, we prove the main lemma.

Remarks 1.2 (1) The main lemma is more transparent than its topological counterpart in [Ni].

(2) After proving the main lemma, enquiries with algebraist colleagues revealed that a more general lemma has already been proved by Colliot-Thélène and Ojanguren (see proposition 1.3 in [C-O]). Our proof is more geometric but less general.

(3) When the total space of C is nonsingular, it is known that (see [H-N] Proposition 1.4 or [Ne] Theorem 2) β is zero only if α is zero. This follows from theorem 1.1 by taking $\tau = 1$, though theorem 1.1 makes a stronger statement even in this case.

2 Reduction to Main Lemma

For basic definitions about conic bundles, see for example Newstead [Ne].

It is clearly enough to prove the theorem for each connected component of X , so we will assume X to be connected. If $Y = \cup_i Y_i$ are the connected components of Y , then by replacing (X, Y) by $(X - \cup_{j \neq i} Y_j, Y_i)$ we are reduced to the case where Y also is connected. Hence we can assume that both X and Y are irreducible.

Let the conic bundle $C \rightarrow X$ be defined via a rank 3 vector bundle E on X , together with quadratic form q on E with values in some line bundle L on X . The quadratic form q is of rank 3 on $X - Y$ because we have a non-degenerate conic over $X - Y$, and q has rank 2 on Y as we have pairs of distinct lines over geometric points of Y .

Let $U = \text{Spec} R$ be an affine open subscheme of X which intersects Y , such that E and L are trivial on U , and $Y \cap U$ is defined by a principal ideal (π) in R . By injectivity of $H^1(Y, \mu_2) \rightarrow H^1(Y \cap U, \mu_2)$, it is enough to prove the theorem for $(U, Y \cap U)$ in place of (X, Y) . Hence we can assume that the conic bundle is defined by an explicit quadratic form $x^2 - ay^2 - bz^2$ on the ring R , where a is a unit in R , while $b \in (\pi)$ vanishes over Y .

Let η be the generic point of Y , and let A be the discrete valuation ring $\mathcal{O}_{X, \eta}$. Let k be the function field of Y . The morphism $\text{Spec}(k) \rightarrow Y$ induces an injective homomorphism $H^1(Y, \mu_2) \rightarrow H^1(k, \mu_2)$. Hence the theorem follows from the following lemma.

Lemma 2.1 (Main Lemma) *Let F be a field of characteristic $\neq 2$. Let A be a discrete valuation ring which is the local ring at the generic point of a smooth divisor in a smooth F -variety. Let K be the quotient field of A , let k be the residue field, and let $\nu : A - \{0\} \rightarrow \mathbb{Z}$ be the discrete valuation. Let $x^2 - ay^2 - bz^2$ be*

a quadratic form on A , with a a unit in A , and $b \neq 0$. Let $(a, b) \in H^2(K, \mu_2)$ be the Brauer class (Hilbert symbol) of the quadratic form on K . Let $\bar{a} \in k - \{0\}$ be the residue class of a , and let $\chi(\bar{a}) \in H^1(k, \mu_2)$ be the class of the two sheeted etale cover $k(\bar{a}^{1/2})/k$ (Kummer character). Then under the Gysin homomorphism $H^2(K, \mu_2) \rightarrow H^1(k, \mu_2)$, we have (in additive notation)

$$(a, b) \mapsto \nu(b)\chi(\bar{a})$$

3 Proof of the Main Lemma

In the course of the proof below, we use the following elementary facts which can be found for example in the textbook of Milne [M]. If S is a field or more generally a Henselian local ring, then $Br(S) \rightarrow H^2(B, \underline{GL}_1)$ is an isomorphism, and provided the characteristic of the residue field is $\neq 2$, the homomorphism $H^2(S, \mu_2) \rightarrow H^2(B, \underline{GL}_1)$ is injective. Moreover the etale cohomology of S with coefficients in a smooth representable sheaf (for example μ_2 or \underline{PGL}_2) is isomorphic by restriction to the corresponding etale cohomology of the residue field of S .

We need two more fact, which are contained in the following two remarks.

Remark 3.1 Let A be a henselian local ring and B/A be a 2-sheeted finite etale cover. If the image of $\gamma \in H^2(A, \mu_2)$ is zero under the composite

$$H^2(A, \mu_2) \rightarrow H^2(B, \mu_2) \rightarrow H^2(L, \mu_2)$$

where L is the quotient field of B , then γ lies in the image of the canonical (connecting) set map $H^1(A, \underline{PGL}_2) \rightarrow H^2(A, \mu_2)$ for the following reason. Any generic section of a Brauer-Severi variety on $Spec(B)$ extends to a global section by the valuative criterion of properness and so the map $H^2(B, \mu_2) \rightarrow H^2(L, \mu_2)$ is injective, hence γ maps to zero under $H^2(A, \mu_2) \rightarrow H^2(B, \mu_2)$. Hence γ is represented by an element (factor set) of the group cohomology $H^2(Gal(B/A), \mu_2)$. As $Gal(B/A)$ is of order 2, the factor set γ defines an Azumaya algebra of rank 2, showing γ comes from $H^1(R, \underline{PGL}_2)$.

Remark 3.2 Let K be a field of characteristic $\neq 2$, let $a, b \in K - \{0\}$ such that a is not a square in K , and let $L = K[t]/(t^2 - a)$. As the conic in P_K^2 defined by $x^2 - ay^2 - bz^2 = 0$ has an L -rational point, its Brauer class (a, b) is an element of the group cohomology set $H^1(Gal(L/K), PGL_2(L))$. Let $Gal(L/K) = \{1, \sigma\}$. A 1-cocycle for $Gal(L/K)$ with coefficients $PGL_2(L)$ therefore consists of an element $g \in PGL_2(L)$ such that $g\sigma(g) = I$ in PGL_2 . (The corresponding ‘crossed homomorphism’ $Gal(L/K) \rightarrow PGL_2(L)$ is defined by $\sigma \mapsto g$.) If $g' \in GL_2(L)$ is an arbitrary lift of g , then there must exist some $c \neq 0$ in L such that $g'\sigma(g') = cI$ in $GL_2(L)$ (which implies $c \in K$). In particular it can be seen by using stereographic projection from $(\sqrt{a}, 1, 0) \in P_K^2(L)$ that (a, b) is represented in $H^1(Gal(L/K), PGL_2(L))$ by

the 1-cocycle defined by

$$h = \begin{pmatrix} & b \\ 1 & \end{pmatrix}$$

We now prove the main lemma.

Proof If b is a unit in A (that is, $\nu(b) = 0$), then the P^1 bundle given by $x^2 - ay^2 - bz^2$ is defined over all of $\text{Spec}(A)$, and so by exactness of the Gysin sequence

$$H^2(A, \mu_2) \rightarrow H^2(K, \mu_2) \rightarrow H^1(k, \mu_2)$$

the image of $(a, b) \in H^2(K, \mu_2)$ in $H^1(k, \mu_2)$ is also zero. So we now assume $\bar{b} = 0$ (which is anyway the case which interests us). If $\nu(b) = 2n$ is even then as $\chi(\bar{a})$ is 2-torsion we have $\nu(b)\chi(\bar{a}) = 0$. Making the change of variable $z' = \pi^n z$ over K , we see that $(a, b) = (a, b') \in H^2(K, \mu_2)$ where $b' = b/\pi^{2n}$. As $\nu(b') = 0$, the argument above now completes the proof when $\nu(b)$ is even. When $\nu(b) = 2n + 1$ is odd, the same change of variables enables us to reduce to the case where $\nu(b) = 1$. Hence we assume from now on that $\nu(b) = 1$.

Note that passing to the completion of A does not affect the residue field and gives a commutative square of the Gysins. Hence it is enough to show the conclusion of the lemma assuming A is complete. If $\bar{a} \in k$ is a square then (by the Henselian property of a complete local ring) a will be a square in A and hence in K , and conversely (this uses the discrete valuation) if $a \in A$ is a square in K then it is a square in A and hence \bar{a} is a square in k . The conic defined in P_K^2 by $x^2 - ay^2 - bz^2$ will have a K rational point $(a_1, 1, 0)$ whenever a_1 is a squareroot of a in K , showing its Brauer class over K is zero. Hence if $\chi(\bar{a}) = 0$ then $(a, b) = 0$, and so the lemma holds. Hence we now assume that \bar{a} is not a square in k , and therefore a is not a square in K or A . (As moreover $\nu(b) = 1$, it follows by remark 1.3 (3) that $(a, b) \neq 0$, though this will also follow from the argument below.)

Let $B = A[t]/(t - a^2)$ which is a 2-sheeted finite etale cover of A , and let L and l be respectively the quotient field and the residue field of B . Consider the commutative diagram where the horizontal maps are the two Gysins.

$$\begin{array}{ccc} H^2(K, \mu_2) & \rightarrow & H^1(k, \mu_2) \\ \downarrow & & \downarrow \\ H^2(L, \mu_2) & \rightarrow & H^1(l, \mu_2) \end{array}$$

The kernel of $H^1(k, \mu_2) \rightarrow H^1(l, \mu_2)$ is of order two, generated by $\chi(\bar{a})$. As a has a squareroot in L , $(a, b) \in Br(K)$ maps to zero in $Br(L)$. Hence to show that $(a, b) \mapsto \chi(\bar{a})$, it is enough to show that the image of (a, b) in $H^1(k, \mu_2)$ is nonzero. By exactness of the Gysin sequence, this will follow if we show that (a, b) does not lie in the image of $H^2(A, \mu_2) \rightarrow H^2(K, \mu_2)$.

Suppose (a, b) was the image of $\gamma \in H^2(A, \mu_2)$. As (a, b) restricts to zero over L , it follows from remark 3.1 that γ lies in the image of some $\gamma' \in H^1(A, \underline{PGL}_2)$ under $H^1(A, \underline{PGL}_2) \rightarrow H^2(A, \mu_2)$. As γ' splits over B , it can be regarded as an element

of the group cohomology set $H^1(\text{Gal}(B/A), \text{PGL}_2(B))$, and hence as in remark 3.2, its 1-cocycle is represented by an element $g \in \text{GL}_2(B)$ such that

$$g\sigma(g) = eI$$

where $\text{Gal}(B/A) = \{1, \sigma\}$ and e is a unit in A .

On the other hand, the group cohomology class of $(a, b) \in H^1(\text{Gal}(L/K), \text{PGL}_2(L))$ can be represented by the matrix h given by remark 3.2. If the cohomology class of (g) were to map to the cohomology class of (h) in $H^1(\text{Gal}(L/K), \text{PGL}_2(L))$, then by definition of group cohomology there would exist elements $M \in \text{GL}_2(L)$ and $0 \neq c \in L$ such that

$$h = cMg\sigma(M^{-1})$$

Applying σ , this would give

$$h = \sigma(h) = \sigma(c)\sigma(M)\sigma(g)M^{-1}$$

Multiplying the two equations and using $h^2 = bI$ and $g\sigma(g) = eI$, we would get

$$b/e = \text{Norm}(c)$$

But this is impossible as on one hand $e \in A$ is a unit while $b \in A$ has valuation $\nu(b) = 1$ so $\nu(b/e) = 1$, and on the other hand ν takes only even values on norms.

Hence the image of (a, b) in $H^1(k, \mu_2)$ is nonzero. This completes the proof of the main lemma and hence that of the theorem.

References

- [C-O] Colliot-Thélène J. L. and Ojanguren M. : ‘Variétés unirationnelles non rationnelles: au-delà de l’exemple d’Artin et Mumford’, *Inventiones Math.* 97 (1989) 141-158.
- [H-N] Hirschowitz, A. and Narasimhan, M. S. : ‘Fibrés de ’t Hooft speciaux et applications’, in *Enumerative geometry and classical algebraic geometry*, Progress in Mathematics 24, Birkhauser, 1982.
- [M] Milne, J. S. : *Étale Cohomology*, Princeton Univ. Press, 1980.
- [Ne] Newstead, P. E. : ‘Comparison theorems for conic bundles’, *Math. Proc. Cambridge Philos. Soc.* 90 (1981) 21-31.
- [Ni] Nitsure, N. : ‘Topology of conic bundles’, *J. London Math. Soc.* (2) 35 (1987) 18-28.

School of Mathematics, Tata Institute of Fundamental Research, Homi Bhabha Road, Mumbai 400 005. e-mail: nitsure@tifrvax.tifr.res.in