



Locally unknotted spines of Heegaard splittings

JESSE JOHNSON

Abstract We show that under reasonable conditions, the spines of the handlebodies of a strongly irreducible Heegaard splitting will intersect a closed ball in a graph which is isotopic into the boundary of the ball. This is in some sense a generalization of the results by Scharlemann on how a strongly irreducible Heegaard splitting surface can intersect a ball.

AMS Classification 57M27; 57Q10

Keywords Heegaard splitting, sweep-out, locally unknotted spine

1 Introduction

The goal of this paper is to understand how a strongly irreducible Heegaard splitting of a manifold M intersects an arbitrary closed ball in $B \subset M$. Intuitively, a strongly irreducible splitting is as efficient as possible so the intersection should be very simple.

This problem was first studied by Scharlemann [4], who showed that under certain conditions, the splitting surface will intersect B as the boundary of a neighborhood of a graph on n points which is isotopic into ∂B .

We will look at how the spine of a Heegaard splitting intersects B . In particular, we will show that under reasonable conditions, the pieces of the spine inside the ball will be unknotted, i.e. simultaneously isotopic into the boundary of the ball.

We will begin by generalizing the notion of a tangle. For a graph T , let ∂T be the set of valence-one vertices.

Definition 1.1 Let B be a closed ball. A *graph tangle* is a graph T (not necessarily connected) properly embedded in B , i.e. such that any valence-one vertices are in the boundary of B .

Definition 1.2 A graph tangle T is *unknotted* if there is an embedding of $T \times [0, 1]$ such that $T \times \{0\}$ maps to T and $(T \times \{1\}) \cup (\partial T \times [0, 1])$ is embedded in ∂B . In other words, an unknotted graph tangle can be isotoped into the boundary of B .

Definition 1.3 Let G be a graph and let $P \subset G$ be a finite collection of points, disjoint from the vertices of G , which separate G . A component of $G \setminus P$ will be called a *cut graph* of G .

Let M be a compact 3-manifold. Let B be a ball embedded in M and let K be a graph embedded in M such that $B \cap K$ is non-empty.

Definition 1.4 A *trivial cone* of K with respect to B is a cut graph C of K (usually a cone on two or more points) and an embedding of $C \times I$ such that $C \times \{0\} = C$, $(C \times \{1\}) \cup (\partial C \times I) \subset \partial B$, the interior of $C \times I$ is disjoint from K and near $(C \times \{1\}) \cup (\partial C \times I)$, $C \times I$ lies outside of B . (Here, $I = [0, 1]$.)

Definition 1.5 A graph, K , (not necessarily connected) in a manifold, M , is *locally unknotted* if for any closed ball B such that there are no trivial cones with respect to B , $K \cap B$ is unknotted in B . Otherwise, K is *locally knotted*.

The requirement that there be no trivial cones is necessary because of the following example: Let K be a graph embedded in M . Let B be a small




Figure 1: A graph which is knotted in a ball because of a trivial cone

ball which intersects an edge of K in an unknotted arc α . Isotope α in B to

form the shape shown in Figure 1. Let B' be the smaller ball, indicated in the Figure. There are two parallel sub-arcs of α in B' , each of which is knotted. The graph K may still be locally unknotted because for this choice of B' , there is a trivial cone.

Definition 1.6 Let M be a compact, connected, closed, orientable 3-manifold. Let Σ be a compact, connected, closed, orientable, two-sided surface in M such that $M \setminus \Sigma$ consists of two components, X_1 and X_2 . Assume $H_1 = X_1 \cup \Sigma$ and $H_2 = X_2 \cup \Sigma$ are handlebodies. Then the ordered triple (Σ, H_1, H_2) is a *Heegaard splitting* for M .

The splitting surface Σ defines the handlebodies H_1 and H_2 up to labeling so we will often refer to the Heegaard surface Σ to indicate the Heegaard splitting (Σ, H_1, H_2) .

A Heegaard splitting is *stabilized* if there are properly embedded, essential disks D_1, D_2 in H_1, H_2 respectively such that ∂D_1 and ∂D_2 intersect transversely in a single point in Σ . A splitting is *reducible* (*weakly reducible*) if there are properly embedded, essential disks in H_1, H_2 , respectively, whose boundaries coincide (are disjoint.)

A *spine* for a handlebody H_1 is a graph K_1 embedded in H_1 such that $H_1 \setminus K_1 \cong \partial H_1 \times (0, 1]$. For a Heegaard splitting (Σ, H_1, H_2) , a spine for Σ is the disjoint union $K = K_1 \cup K_2$ where K_1 is a spine for H_1 and K_2 is a spine for H_2 .

We will prove the following:

Main Theorem *Let (Σ, H_1, H_2) be a strongly irreducible Heegaard splitting for a manifold M . Then every spine of Σ is locally unknotted.*

In Section 2, we define sweep-outs and introduce the Rubinstein-Scharlemann graphic. This is the main tool used in Section 3 to prove Proposition 3.1. The Proposition states that if a Heegaard splitting is strongly irreducible then some level surface of any sweep-out intersects a given ball in trivial pieces. This is essentially a modification of the result in [3] that given any two sweep-outs, there is a pair of level surfaces such that each loop of intersection is either trivial in both surfaces or non-trivial in both surfaces. In our case, ∂B contains only trivial loops so the loops must all be trivial in the level surface of f as well. The proof of the Main Theorem is given in the final section.

To indicate the significance of this theorem, we note that if Σ is stabilized then there is a spine of Σ which is locally knotted. To see this, let (Σ, H_1, H_2) be




Figure 2: A locally knotted spine of a stabilized Heegaard splitting

a stabilization of a strongly irreducible Heegaard splitting (Σ', H'_1, H'_2) . Let K be a spine of H'_1 with a single vertex. Because Σ' is strongly irreducible, any edge e of K is not isotopic into a ball. Form a spine for H_1 by adding a trivial loop e' to K . Slide e' along e to create a formation as in Figure 2 and let B be the ball indicated by the dotted line. The two arcs are knotted in B and there are no trivial cones (because e is not isotopic into a ball) so H_1 is locally knotted.

This research was supported by NSF VIGRE grant 0135345.

2 The Rubinstein-Scharlemann graphic

Let M be a compact, connected, closed, orientable 3-manifold and (Σ, H_1, H_2) a Heegaard splitting for M . Let K_i be a spine for H_i . Then $H_i \setminus K_i$ is homeomorphic to $\Sigma \times (0, 1]$. Let $h : (H_i \setminus K_i) \rightarrow \Sigma \times (0, 1] : x \mapsto (p_x, l_x)$ be a homeomorphism and let $f : H_i \rightarrow I$ such that $f(x) = l_x$ when $x \notin K_i$ and $f(x) = 0$ when $x \in K_i$. If we combine the two functions (one for each handlebody,) we get a function on all of M .

Definition 2.1 A *sweep-out* is a smooth function $f : M \rightarrow [-1, 1]$ such that $f^{-1}(-1)$ is a spine of H_1 , $f^{-1}(1)$ is a spine of H_2 and $f^{-1}(t)$ is a surface isotopic to Σ for each $t \in (-1, 1)$.

By choosing a smooth embedding for each K_i and a smooth map h , the above construction produces a sweep-out for any Heegaard splitting of M .

Definition 2.2 Let f and g be sweep-outs and define $f_t = f^{-1}(t)$ and $g_t = g|_{f_t}$. We will say that f and g are in *general position* if g_t is a Morse

function on f_t for all but finitely many t , and g_t is a near-Morse function for the remaining values. (A function is *near-Morse* if there is a single degenerate critical point or there are two non-degenerate critical points at the same level.)

Note that for $t \in (-1, 1)$, f_t is a surface embedded in M , while g_t is a function on f_t .

Let f and g be sweep-outs in general position. For each $t \in [-1, 1]$, let s_t be a critical value of the function g_t . Rubinstein and Scharlemann [3] have shown that the points $(t, s_t) \in [-1, 1] \times [-1, 1]$ form a graph $G \subset \mathbf{R}^2$ with vertices of valence 2 and 4 in the interior and valence 1 and 2 on the edges of




Figure 3: A possible Rubinstein-Scharlemann graphic

$[-1, 1] \times [-1, 1]$. We will call this graph the *Rubinstein-Scharlemann graphic*.

The valence-2 vertices in the interior arise when g_t is a near-Morse function with a degenerate critical point. As t passes the vertex, a central singularity and a saddle singularity in f_t come together and cancel.

At a valence-4 vertex, (t, s_t) , g_t has two critical points at the same level. As t passes the vertex, one of the critical points rises past the other. The two critical points may be on the same component of $g_t^{-1}(s_t)$ or different components.

If points (t, s) and (t', s') are in the same component of $([-1, 1] \times [-1, 1]) \setminus G$ then there is an isotopy of the surface f_t onto the surface $f_{t'}$ taking the collection of loops $g_t^{-1}(s)$ to the loops $g_{t'}^{-1}(s')$. In other words, the topological properties of the loops in the level set depend only on the component c , not the choice of a point within c .

In the following section, we will replace the sweep-out g with a sweep-out of a ball $B \subset M$, i.e. a map $g : B \rightarrow [0, 1]$ such that $g^{-1}(0)$ is a single point in B and $g^{-1}(s)$ is a sphere parallel to ∂B for every $s \in (0, 1]$.

In this case, the definition of general position is essentially the same. The function g_t is defined on a closed submanifold of f_t . The resulting graphic, as in Figure 3, looks like the bottom half of the Rubinstein-Scharlemann graphic for two sweep-outs.

3 How a sweep-out intersects a closed ball

Let (Σ, H_1, H_2) be a Heegaard splitting of M with spine K . Let $B \subset M$ be a closed ball such that K is transverse to ∂B . Let g be the genus of Σ .

Proposition 3.1 *Suppose for each surface f_t there is an essential curve of f_t lying completely in B . If $g = 1$ then Σ is stabilized and $M = S^3$. If $g \geq 2$ then Σ is weakly reducible.*

Proof Choose a point $x \in B$ away from K . Let N be a regular neighborhood of B and let $g : N \rightarrow [0, 1]$ be a sweep out of N , such that $g^{-1}(0) = \{x\}$. By work of Cerf [2], we can isotope g by an arbitrarily small amount so that g and f are in general position.

In particular, isotope and rescale g so that $B \subset g^{-1}([0, 1]) \subset N$. Let $\{(t_i, s_i)\}$ be the vertices of the Rubinstein-Scharlemann graphic, ordered so that $t_i < t_{i+1}$. (i.e. f_{t_i} has either a degenerate critical point at s_i or two non-degenerate critical points at level s_i .)

The assumption that there is an essential curve of f_t lying inside B is equivalent to assuming that some level curve of g_t is essential in f_t . If g_t is Morse and each level curve is trivial then $f_t \cap B$ is either a collection of disjoint disks or a sphere. In either case, there are no essential loops of f_t lying inside B . Thus for the remainder of the proof, we will assume that for each t , some level curve of g_t is essential in f_t .

Let G be the Rubinstein-Scharlemann graphic in $[-1, 1] \times [0, 1]$. For each component c of $([-1, 1] \times [0, 1]) \setminus G$, let (t_c, s_c) be a point in the interior of c . Let L_c be the collection of loops $g_{t_c}^{-1}(s_c)$. If one of the loops bounds a properly embedded, essential disk in the handlebody $f^{-1}[-1, t_c]$, label the region c with a 1 and if a loop bounds a disk in $f^{-1}[t_c, 1]$, label c with a 2. This label is independent of our choice of (t_c, s_c) within c . Before continuing, we need the following lemma:

Lemma 3.2 *Let (Σ, H_1, H_2) be a Heegaard splitting for M and let S be a 2-sphere embedded in M such that $S \cap \Sigma$ is a non-empty collection of loops. If one or more of the loops is essential in Σ then there is a loop of $S \cap \Sigma$ which bounds a properly embedded, essential disk in either H_1 or H_2 .*

Proof Induct on the number of components of $S \cap \Sigma$. If there is a single component l of $S \cap \Sigma$ then l must be essential. This l splits S into two disks, each of which is a properly embedded, essential disk in one of the handlebodies.

If there are more than one components of $S \cap \Sigma$, let n be the number of loops and assume the result is true whenever there are $n - 1$ loops. Let l be an innermost loop of $S \cap \Sigma$ (one that bounds a disk D in S disjoint from Σ .) If l is essential, then D is properly embedded and essential and we are done.

Otherwise, D is boundary parallel (since handlebodies are irreducible) so we can isotope S to a surface S' such that $S' \cap \Sigma$ is a proper subset of $S \cap \Sigma$. Since $S' \cap \Sigma$ has $n - 1$ components, one of them bounds a properly embedded, essential disk, so the corresponding component of $S \cap \Sigma$ does as well. \square

It follows from the lemma that if a component c of $([-1, 1] \times [0, 1]) \setminus G$ is not labeled then all the loops of L_c are trivial in f_t . If one of the components has both labels 1 and 2, then the Heegaard splitting is weakly reducible, so we can assume each component has at most one label.

If a vertical line in $I \times I$ intersects both a component labeled with a 1 and a component labeled with a 2 then the Heegaard splitting is weakly reducible: Choose t so that the vertical line $\{t\} \times I$ intersects both types of components. Then one level set of g_t will contain a loop bounding a disk in H_1 and another level set will contain a loop bounding a disk in H_2 . The level curves are disjoint so the disks will have disjoint boundaries.

Assume Σ is not weakly irreducible. We will show that Σ must be genus-one and stabilized. For each $t \in I$, if g_t is a Morse function then there is an essential level curve of g_t . So, each vertical line $\{t\} \times I$ must pass through either a vertex of G or one or more labeled components, all with the same label. In the latter case, let l_t be this label.

For t near -1 , f_t approaches the spine K_1 , so l_t must be 1 for small enough t . (Recall, K is transverse to ∂B .) Similarly, l_t must be 2 for t near 1. Thus the label must switch from 1 to 2 somewhere. It can only change at a vertex of G , so let $v = (t_i, s_i)$ be a vertex where $l_{t_i - \epsilon}$ is 1 and $l_{t_i + \epsilon}$ is 2.

The vertical line at t_i passes through only one vertex. Because each vertical line to the left or right of t_i can only pass through one type of component, the vertex v must be a crossing with the regions above and below it unlabeled. This type of crossing is shown in Figure 4.

Lemma 3.3 *Let g be a Morse function on a compact, connected, closed, orientable surface F with critical values c_i . For some i , let $h_1 \in (c_i, c_{i+1})$, $h_2 \in (c_{i+1}, c_{i+2})$ and $h_3 \in (c_{i+2}, c_{i+3})$. If $g^{-1}(h_1) \cup g^{-1}(h_3)$ consists of trivial loops and $g^{-1}(h_2)$ contains non-trivial loops then F is a torus.*




Figure 4: A crossing at which l_c switches, without implying that Σ is weakly reducible

Proof The level set $g^{-1}(h_2)$ is the result of adding a band to some loop $a_1 \subset g^{-1}(h_1)$. By assumption, a_1 must be trivial so let $D_1 \subset F$ be the disk whose boundary is a_1 . Let $D_2 \subset F$ be the 1-handle which is attached to D_1 .

The level set $g^{-1}(h_3)$ is the result of adding a second 1-handle. Because all the loops at this level are trivial, the band must be attached to $\partial(D_1 \cup D_2)$. Let D_3 be this 1-handle and let $a_3 = \partial(D_1 \cup D_2 \cup D_3)$. Then a_3 must bound a disk D_4 because a_3 is isotopic to a loop in $g^{-1}(h_3)$.

This D_4 must be disjoint from the other handles so the four disks suggest a handle decomposition of F . Therefore the Euler characteristic of F is 0 and F is a torus. \square

We can now complete the proof of Proposition 3.1. If l_c changes at a crossing as in Figure 4 then the Morse function $g_{t_i-\epsilon}$ satisfies the conditions of Lemma 3.3, implying Σ is a torus. This proves Proposition 3.1 when the genus of Σ is greater than one. The only case that remains is when the Heegaard splitting has genus one.

Let $g_1 = g_{t_i-\epsilon}$ and let $g_2 = g_{t_i+\epsilon}$. The only difference between the two functions is the order of the 2-handles that create and destroy the essential loops. One can check that the loops created by adding the 1-handles in different orders will intersect in a single point and bound disks in H_1, H_2 , respectively.

Thus if Σ is not weakly reducible then Σ is a stabilized, genus-one Heegaard splitting (of S^3). \square

Corollary 3.4 (Waldhausen's Theorem) *Every positive-genus Heegaard splitting of S^3 is stabilized.*

Proof Let (Σ, H_1, H_2) be a Heegaard splitting of S^3 with genus g . Let f be a sweep-out for Σ . Let N be a small ball in S^3 such that $f^{-1}(t) \cap N$ is either

a point, a disk or empty for all t . Then $B = S^3 \setminus N$ is a ball and every level set of f intersects B non-trivially.

If $g = 1$ then Σ is stabilized and we are done. If $g \geq 2$ then Σ is weakly reducible. The main theorem in [1] states that if a Heegaard splitting is weakly reducible then it is reducible or the ambient manifold is Haken. Because S^3 contains no incompressible surfaces, this implies that Σ is reducible.

Thus Σ is a connect sum of Heegaard splittings of S^3 with strictly lower genus. By induction, Σ is a connect sum of g splittings, each with genus 1. Because each of these splittings is stabilized, their connect sum Σ is stabilized. \square

Note that the proof of Corollary 3.4 is a modification of Theorem 5.11 in [3].

4 Proof of the Main Theorem

We will begin by stating three lemmas which will be necessary in the final proof. A proof of the first lemma will be left to the reader.

Lemma 4.1 *Let T be a graph tangle in a ball, B , and let $D \subset B$ be a properly embedded disk, disjoint from T , cutting B into two balls B_1 and B_2 . Assume that $T \cap B_1$ is unknotted in B_1 . Then T is unknotted in B if and only if $T \cap B_2$ is unknotted in B_2 .*

Let Σ be a strongly irreducible Heegaard splitting and let K and B be as in Section 3. Assume that K has no trivial cones with respect to B .

Lemma 4.2 *If Σ is strongly irreducible and $B \subset M$ is a closed ball then there is an immersion of $K \times I$ such that $K \times [0, 1)$ is embedded, $K \times \{0\} = K$ and $K \times \{1\}$ is disjoint from B .*

Proof Let f be a sweep-out for Σ . By Proposition 3.1, there must be a value $t \in (-1, 1)$ such that $B \cap f_t$ does not contain any essential loops of f_t . Let L be the collection of loops $f_t \cap \partial B$. These loops are pairwise disjoint and each loop $\alpha \in L$ bounds a disk $D_\alpha \subset f_t$.

There is an embedding of $K_1 \times [0, 1]$ in $f^{-1}[-1, t]$ such that $K_1 \times \{0\} = K_1$ and $K_1 \times \{1\} \subset f_t \setminus \bigcup D_\alpha$. Then $K_1 \times \{1\}$ is disjoint from L so $K_1 \times \{1\}$ is either in B or disjoint from B . If it were inside B , then K_1 would be isotopic

into B (a contradiction) so it must be disjoint from B . The same construction for K_2 finishes the proof.

Note that in general, $K_1 \times \{1\} \cap K_2 \times \{1\}$ will be non-empty. The statement of the lemma requires only that the interior of $K \times I$ be embedded. \square

Let $K \times [0, 1]$ be immersed in M with embedded interior such that $K \times \{0\} = K$ and $K \times \{1\}$ is disjoint from B , as in Lemma 4.2. We will write $K \times I$ for the image of this immersion. Let $L = (K \times I) \cap \partial B$.

Let e be an edge of K . Then $L \cap (e \times I)$ consists of a number of arcs and loops. Because $K \times \{1\}$ is disjoint from B , each arc in $e \times I$ must have both its endpoints in $K \times \{0\}$ or in $v \times (0, 1)$ where v is an endpoint of e . Let α be such an arc.

If α lies in $e \times [0, 1]$ and has both its endpoints on the same $v \times (0, 1)$ for some v , then the endpoints of α bound a sub-arc β of $v \times (0, 1)$. The loop $\alpha \cup \beta$ bounds a disk $D \subset e \times [0, 1]$. Sliding β across D gives an isotopy of $v \times (0, 1)$ which extends to $K \times (0, 1)$. This isotopy reduces the number of intersections of $v \times [0, 1]$ with ∂B . Thus by a finite number of such isotopies, we can eliminate all arcs with both endpoints on the same vertical edge $v \times (0, 1)$.

Lemma 4.3 *Let l be a component of L which intersects $K \times \{0\}$. Then l separates $K \times I$ into a component homeomorphic to $K \times I$ and a component homeomorphic to $l \times I$.*

Proof Consider a projection ρ of L into $K \times \{0\}$. After eliminating arcs in L with both endpoints on the same edge $v \times I$, we can choose a projection ρ so that ρ is one-to-one on each arc of L . We will prove that ρ is then one-to-one on the whole component l . Assume for contradiction this map sends points $x, y \in l$ to the same point in $K \times \{0\}$ and assume y is below x (i.e. farther from $K \times \{0\}$.)

Since l is connected, there is a path $\alpha : [0, 1] \rightarrow l$ with $\alpha(0) = x$ and $\alpha(1) = y$. We would like to construct a path $\alpha' : [0, 1] \rightarrow l$ such that $\rho \circ \alpha(t) = \rho \circ \alpha'(t)$ for every t and $\alpha'(0) = y$. This is, in some sense, a lift of the projection of α , starting at a different base point. We will show that for each t , $\alpha'(t)$ is below $\alpha(t)$.

The edges of $v \times I$ of $K \times I$ cut α into a number of arcs coinciding with the arcs of L . The first arc can't end on $K \times \{0\}$ because then α would end. It can't end at y because ρ is one-to-one on each arc of l . Thus the first arc must pass from x to some $v \times I$.

The point y also sits on an arc of l . This arc extends in the same direction as the first arc of α . It can't end on $K \times \{0\}$ because it is below another arc of L which extends to $v \times I$. It cannot end on $K \times \{1\}$ because L is disjoint from $K \times \{1\}$. Thus the arc ends on $v \times I$, below the first arc of α . We can define α' as desired, strictly below the first arc of α .

Continue in this fashion for each arc in the image of α , until α reaches y . Now, $\alpha'(1)$ is a point z below y such that $\rho(z) = \rho(y) = \rho(x)$. Continuing the construction, we can find another point below z , and so on, implying that $\rho^{-1}\rho(x)$ contains an infinite number of points. But this is impossible because L is a finite graph. Thus ρ must be one-to-one.

For each point $x \in l$, consider the vertical arc a_x from x to $K \times \{0\}$. Because ρ is one-to-one, the union of these arcs is homeomorphic to $l \times I$. Its complement is homeomorphic to $K \times I$. □

We are now ready to prove the main theorem. Isotope $K \times (0, 1)$ so as to minimize the number of loops in L and the number of intersections with vertical edges $v \times I$. Each component of $K \cap B$ contains one or more components of L . If there are no loops, then each $l \times I$ component lies entirely in B so by Lemma 4.3, each l cuts off a piece of $K \times I$ homeomorphic $l \times I$. This defines an isotopy of a component of $K \cap B$ into ∂B so $K \cap B$ is unknotted. Otherwise, if there are loops in L , we will show that there is always a disk properly embedded in B which cuts off an unknotted tangle, allowing us to use Lemma 4.1.

Let β be a loop of L . Then β bounds a disk, $D_1 \subset (K \times [0, 1])$. Assume β is an inner-most loop, i.e. assume $D_1 \cap \partial B = \partial D_1$. There are two cases to consider: either D_1 lies entirely in B or the interior of D_1 lies outside of B .

Assume the interior of D_1 is disjoint from B . The loop β separates ∂B into two disks. Because M allows a strongly irreducible Heegaard splitting, M is irreducible, so $D_1 \cup \partial B$ separates M into two balls and a punctured M . Let D_2 be the disk of $\partial B \setminus \beta$ such that $D_1 \cup D_2$ bounds a ball, B' outside B .

The graph $K \times \{1\}$ is disjoint from $\partial B'$ so each component of $K \times \{1\}$ must lie outside of B' . (If it were inside B' then $K \times I$ would suggest an isotopy bringing one of the spines into B' .)

If $K \times I$ intersects D_2 , then let l be a component of $D_2 \cap K \times I \subset L$. By Lemma 4.3, l cuts off a piece of $K \times I$ homeomorphic to $l \times I$. This piece is outside of B so $l \times I$ defines a trivial cone. Thus there are no arc-intersections and $K \times \{0\}$ lies outside of B' . Any components of $K \times I$ in B' can be

isotoped out of B' and then D_1 can be isotoped into B , reducing the number of components of L .

So, if L is minimized, we can assume no loops bound disks whose interiors are disjoint from B . Let \mathbf{D} be the collection of innermost disks in L . These separate B into a number of balls. Let $D \in \mathbf{D}$ be an outermost disk in B , cutting off a ball B' which does not contain any disks of \mathbf{D} . Let B'' be the complement of B' . Note that a trivial cone of K with respect to B' or B'' is also a trivial cone with respect to B .

If there are no arcs of K in B' , then we can isotope all the pieces of $K \times I$ out of B' and then isotope off D . Otherwise, let $L' = (K \times I) \cap \partial B'$ and let β be an innermost loop of L' cutting off a disk D' .

The disk D' cannot lie in B' because D was assumed to be an outermost disk. Thus D' must lie outside B' . By applying the argument above, to B' instead of B , it is clear that D' can be isotoped into B' without affecting K . So, if the number of loops of intersection is minimized, there can be no loops in L' .

Thus $K \cap B'$ is an unknotted tangle in B' . By Lemma 4.1, $K \cap B$ is unknotted if and only if $K \cap B''$ is unknotted. Again minimize the number of loops of intersection for this new tangle. The number of components of $(K \times I) \cap (B \setminus B')$ is strictly less than the number of components of $(K \times I) \cap B$ so the process will eventually terminate, proving the theorem. \square

References

- [1] **A J Casson, C M Gordon**, *Reducing Heegaard splittings*, Topology Appl. 27 (1987) 275–283 MR918537
- [2] **J Cerf**, *Sur les difféomorphismes de la sphère de dimension trois ($\Gamma_4 = 0$)*, Lecture Notes in Mathematics, No. 53, Springer-Verlag, Berlin (1968) MR0229250
- [3] **H Rubinstein, M Scharlemann**, *Comparing Heegaard splittings of non-Haken 3-manifolds*, Topology 35 (1996) 1005–1026 MR1404921
- [4] **M Scharlemann**, *Local detection of strongly irreducible Heegaard splittings*, Topology Appl. 90 (1998) 135–147 MR1648310

Mathematics Department, University of California
Davis, CA 95616, USA

Email: jjohnson@math.ucdavis.edu

Received: 27 January 2005 Revised: 9 June 2005